


# Adult Cardiac Arrest Algorithm – 2018 Update



## CPR Quality

- Push hard (at least 2 inches [5 cm]) and fast (100–120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 30:2 compression-ventilation ratio.
- Quantitative waveform capnography
  - If  $\text{PETCO}_2 < 10 \text{ mm Hg}$ , attempt to improve CPR quality.
- Intra-arterial pressure
  - If relaxation phase (diastolic) pressure  $< 20 \text{ mm Hg}$ , attempt to improve CPR quality.

## Shock Energy for Defibrillation

- Biphasic:** Manufacturer recommendation (eg, initial dose of 120–200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered.
- Monophasic:** 360 J

## Drug Therapy

- Epinephrine IV/IO dose:** 1 mg every 3–5 minutes
- Amiodarone IV/IO dose:** First dose: 300 mg bolus. Second dose: 150 mg.  
**-OR-**  
**Lidocaine IV/IO dose:** First dose: 1–1.5 mg/kg. Second dose: 0.5–0.75 mg/kg.

## Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

## Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Abrupt sustained increase in  $\text{PETCO}_2$  (typically  $\geq 40 \text{ mm Hg}$ )
- Spontaneous arterial pressure waves with intra-arterial monitoring

## Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary