

Surgical management of Inflammatory bowel disease

- indication of surgical intervention
- understand the risk involved
- option and rationale of surgical treatment



Multi-disciplinary care

- ◉ Named personnel comprising
 - > gastroenterologists,
 - > colorectal surgeons
 - > clinical nurse specialists,
 - > dietician,
 - > pathologist
 - > GI radiologist
 - > pharmacist,
- ◉ Access to
 - > a psychologist/counsellor, rheumatologist, ophthalmologist,
 - > dermatologist, obstetrician, nutrition support team, a paediatric gastroenterologist
 - > gastroenterology clinical network, general practise



Surgical management of Ulcerative colitis





Risk of Surgery / UC

- ⦿ 20 – 30 % of patients will require surgery
- ⦿ 5-10 % present with acute sever colitis
- ⦿ 30 % of sever case will require emergency surgery
- ⦿ After acute severe ulcerative colitis 50% with incomplete remission with steroids will require colectomy within 1 year.



Acute Severe Colitis

- 6 bloody stools/day
- Abdominal tenderness
- signs of systemic Toxicity (HR>90, T>37.8, Hb#10.5 or ESR>30)
- Anemia
- Fulminant colitis (stool > 10 / day , Anemia requiring transfusion , signs of systemic toxicity , abdominal distension, tenderness , fever and leukocytosis.

Truelove and Witts' criteria , Br Med J. 1955

Sands BE J Gastrointestinal Surg. 2008

- Colectomy rate about 30 %
- Rate of Colectomy did not change in last 40 years

Turner D. et al Gastroenterol Hepatol 2007



	Mild	Moderate	Severe
1. Number of evacuations/day	≤4	5	≥6
2. Bright-red blood in stool	-	+	++
3. Temperature (°C)	Normal	Intermediate values	Average temperature at night >37.5 °C or >37.8° C in 2 days within 4 days
4. Pulse (bpm)	Normal	Intermediate	>90 bpm
5. Hemoglobin(g/dL)	>10	Intermediate	≤10.5
6. *HSS (mm, 1st hour)	≤30	Intermediate	>30

*HSS : Hemocritinization speed

FIGURE 2. Classification of nonspecific ulcerative colitis (UC) according to severity of acute episode (Truelove & Witts⁽⁹³⁾)

Rule of Surgery in Acute Presentation

- ⦿ Perforation
- ⦿ Haemorrhage
- ⦿ Toxic megacolon (diameter >5.5 cm, or caecum >9 cm)
 - > Systemic toxicity
 - > Steroids mask clinical picture.
- ⦿ Failed medical treatment



Colectomy in Acute presentation

- Up to 40 % mortality for **perforation**
- 2-8 % mortality if before perforation



Rule of Surgery in Acute Presentation . Cont.

- ⊙ GI team care. Surgeon aware.
 - > Routine bloods (CBC/ U&E's / CRP/Albumin)
 - > Regular abdominal exam
 - > AXR
 - > Stool for bacteriology/ C diff / CMV
 - > +/- Flexible sigmoidoscopy
 - > DVT prophylaxis

- ⊙ IV steroids



Rule of Surgery in Acute Presentation . Cont.

- ⊙ A stool frequency of >8/day or CRP >45 mg/l at **3 days** appears to predict the need for surgery in 85% of cases

Travis, S. P. Let al Gut. 38(6):905-910, June 1996.

- ⊙ Intravenous steroids are generally given for up to **5 days**. There is no benefit beyond 7-10 days

Turner D et al Clin Gastroenterol Hepatol 2007;5:103e10.



Rule of Surgery in Acute Presentation . Cont.



- ⦿ Day 1
- ⦿ Day 3 : *Surgery Discussed / Stoma therapist input.*
- ⦿ Day 5
- ⦿ → Consideration of **colectomy** or **rescue** therapy with either intravenous cyclosporine OR Biologic

Surgery in Acute presentation cont.



- ⊙ Proctocolectomy and Ileostomy
 - > High mortality
 - > Permanent stoma
 - > Pelvic dissection / nerve damage / sepsis
- ⊙ Proctocolectomy and Pouch
- ⊙ Subtotal Colectomy and Ileostomy



Subtotal Colectomy and Ileostomy

- ⊙ ~ 3% mortality
- ⊙ ~ 2 -12 % rectal stump blowout.
- ⊙ Close stump / Mucus fistula / SC Stump

Advantage

- > Confirm diagnosis
 - > Off medication
 - > Improve nutritional status
-
- ⊙ *6 months to next stage.*



Options after surgery for acute colitis

- ⊙ Ulcerative colitis / Indeterminate
 - > Completion proctectomy and Pouch
 - > Completion proctectomy and end ileostomy
 - > Completion proctectomy and Continent ileostomy

- ⊙ Crohn's Disease ?



Elective Surgery for Ulcerative Colitis

- Medical Intractability ? Failed medical treatment
 - > MDT
- Chronic disease
 - > Quality of life
 - > Off work / Hospitalization
 - > Never remission / Anemia / Amenorrhea/ ,malnutrition
- Steroid dependence / refractory
- Extra-alimentary manifestation
- Malignancy



Extra intestinal Manifestations

- > Peripheral arthritis
- > Uveitis
- > Iritis

Respond to colectomy

- > Ankylosing spondylitis
- > Sacroilitis
- > Primary sclerosing cholangitis


do not respond to colectomy



Malignancy



- 1 -2 % per year after 10 years
- PSC 9% after 10 years
50 % after 25 years
- Surveillance
- High Grade Dysplasia vs low grade dysplasia
- Pancolitis



Risk of malignancy in UC

- ◉ Pancolitis
- ◉ PSC (primary sclerosing cholangitis)
- ◉ Dysplasia





Options of elective surgery


⊙ Restorative proctocolectomy

- > One or Two stages
- > Reduce steroid to minimum

⊙ Proctocolectomy and end Ileostomy


Restorative proctocolectomy

- Elective
- Off steroids
- One or two stages - w/o ileostomy
- Specialized Units
 - > At least 10 per year BSG 2010 (UK)
- stapled or hand-sewn pouch
- pouch configuration (W, S, J)
- hand-sewn or stapled ileo-anal anastomoses



Life style operation

- The median frequency of defaecation/24
 - > 5 day
 - > 1 night
- Nocturnal seepage
 - > 8% at 1 year
 - > 15 % at 20 years
- Urgency
 - > 5.1% at 1 year
 - > 9.1 % at 15 years





Tekks PP et al Colorectal Dis 2010

- Fecundity reduced by 40 – 50 %

Gorgun E et al Surgery 2004



J or W pouch



McCormick, P. H. et al *Diseases of the Colon & Rectum.*, December 2012.

Risk of malignancy and dysplasia in rectal cuff

- ◉ Low risk / is infrequent

Remzi , Dis Colon Rectum. 2003;

Fazio 1994


- ◉ Cuff surveillance is not necessary
 - > Unless dysplasia and cancer in original sp.

Coul Colorectal Disease. 2007.



Complication after RPC

- ⦿ Pouchitis up to 50 %
 - > Consider CD.
 - > Antibiotics/ Probiotics/ Biologic/ Ciclosporin
- ⦿ Pouch vaginal fistula
 - > Technical
 - > Advancement flaps / redo-pouch
- ⦿ Vitamin B12 and iron deficiency
- ⦿ Infertility
- ⦿ Stricture
- ⦿ Malignancy






You're muted.

Press Ctrl+Shift+M to unmute your microphone.


Complication after RPC

- ⦿ Pouchitis up to 50 %
 - > Consider CD.
 - > Antibiotics/ Probiotics/ Biologic/ Ciclosporin
- ⦿ Pouch vaginal fistula
 - > Technical
 - > Advancement flaps / redo-pouch
- ⦿ Vitamin B12 and iron deficiency
- ⦿ Infertility
- ⦿ Stricture
- ⦿ Malignancy



Pouch failure

- ⊙ 5.9 % at 10 years
 - > Pelvic sepsis
 - > Anastomotic leak
 - > Fistula
 - > Crohn's disease



Fazio *Annals of Surgery* 2013

Surgery of Ulcerative colitis



- ◉ Curative
- ◉ Risk of cancer / Dysplasia
- ◉ Dealing with complication and failure
- ◉ Re-operative / Re-do Surgery
- ◉ Attractive for minimally invasive surgery
- ◉ Controversies remains

Surgery for Crohn's disease

● Indication

- > Stenosis (stricture) causing obstructive symptoms
- > Enterocutaneous or intra-abdominal fistula
- > Intra-abdominal or retroperitoneal abscess
- > Acute or chronic bleeding
- > Free perforation





(Complication)



Surgery CD cont.



- ◎ Segmental resection
- ◎ Avoid wide resection
- ◎ May need stoma
 - > Malnutrition
 - > Immuno-suppression
 - > Intra-abdominal sepsis
- ◎ Risk of malignancy



Strictureplasty





Smoking

- Tobacco abuse as a causative factor in the development of Crohn's disease has been difficult to prove
- Increase the incidence of relapse and failure of maintenance therapy.
- Associated with the severity of disease in a linear dose-response relationship.