


A cartoon illustration of a young boy with brown hair and a blue shirt. He has a wide-eyed, worried expression. A small amount of red blood is dripping from his nose. He is holding his right index finger up to his nose, with a red smudge on the tip. The background is a dark blue gradient with a pattern of light blue circuit lines on the left side.

EPISTAXIS

BY: AHMED SALAHAT &
LEITH SHURIQUIE



CONTENTS:


- 1. Introduction
 - 2. Classification
 - 3. Etiology
 - 4. Treatment
 - 5. Epidemiology
 - 6. History, physical & DDx
 - 7. Complications
 - 8. Prevention
- 

INTRODUCTION

- Epistaxis is the medical term for **nosebleed**.
- It more commonly occurs in **children (ages 2–10)** and **older adults (ages 50–80)**
- It is rarely life threatening but it may cause significant concern, especially among parents of small children.
- Most nosebleeds are **benign, self-limiting**, and **spontaneous**, but some can be **recurrent**.


INTRODUCTION

- The nose has a rich vascular supply, with substantial contributions from the **internal carotid artery (ICA)** and the **external carotid artery (ECA)**.




INTRODUCTION

- The **ECA system**: the **facial** and **internal maxillary** arteries.
 - The **superior labial artery** is one of the terminal branches of the facial artery which supplies the anterior septum through a septal branch.
 - The **internal maxillary artery** enters the pterygomaxillary fossa and divides into 6 branches, some of which are the **greater palatine** and **sphenopalatine arteries**.




INTRODUCTION

- The ICA contributes to the nasal vascularity through the **ophthalmic artery** and its **anterior and posterior ethmoidal branches**.




INTRODUCTION

- **Kiesselbach plexus**, or **Little's area**, is an anastomotic network of vessels located on the anterior cartilaginous septum. It receives blood supply from both the ICA and the ECA. It is where vessels from both the ICA (anterior and posterior ethmoidal arteries) and the ECA (sphenopalatine and branches of the internal maxillary arteries) converge.



INTRODUCTION

- **Woodruff's plexus** is a collection of arteries located in the posteroinferior region of the lateral nasal cavity, formed by anastomoses of the sphenopalatine artery (branch of the maxillary artery) and pharyngeal artery.



CLASSIFICATION

- There are two types of epistaxis depending on their origin: **anterior** and **posterior** epistaxis.

Criteria	Anterior epistaxis	Posterior epistaxis
Clinical features	Bleeding from the nostrils.	*Bleeding through the posterior nasal aperture down the throat (no external signs of bleeding). *Haemoptysis, hematemesis, and/or melena may occur due to swallowing of large amounts of blood.
Relative frequency	90% of cases.	10% of cases.
Peak Incidence	Children < 10 years of age.	Older individuals (> 50 years of age).
Most common site of bleeding	Kiesselbach plexus.	Woodruff plexus.

CLASSIFICATION

EPISTAXIS

ANTERIOR NOSEBLEEDS
ARE MOST COMMON

COMMONLY CAUSED BY
TRAUMA, INCLUDING
NOSE-PICKING

KIESSELBACH'S
PLEXUS


MORE COMMON
IN CHILDREN AND
ADOLESCENTS

WOODRUFF'S PLEXUS

POSTERIOR BLEEDING
IS LESS COMMON

ASSOCIATED WITH
HYPERTENSION AND
ATHEROSCLEROSIS

USUALLY SEEN
IN THE ELDERLY



CLASSIFICATION

- A **posterior** source presents a greater risk of airway compromise, aspiration of blood, and greater difficulty controlling bleeding.

Posterior epistaxis can be a sign of life-threatening hemorrhages!

ETIOLOGY

- In most cases, the exact cause of epistaxis remains unknown (**idiopathic epistaxis**).
- While a single episode of epistaxis usually does not require any investigation, recurrent epistaxis must be investigated for an underlying cause (e.g. a bleeding disorder).
- Causes can be:
 - **Local.**
 - **Systemic.**

ETIOLOGY

- Local Trauma is the most common cause!

LOCAL CAUSES



LOCAL
TRAUMA



ANATOMICAL
IRREGULARITIES



INFLAMMATION



FACIAL
TRAUMA



TOPICAL
NASAL SPRAYS
(incorrect/excessive use)




TUMORS
(rare)

SYSTEMIC CAUSES



HIGH BLOOD
PRESSURE



VASCULAR
MALFORMATIONS



CARDIOVASCULAR
DISEASES



BLEEDING
DISORDERS

TREATMENT

- Management of epistaxis depends on the **severity of the bleeding** and the individual's **accompanying medical conditions**.
- Treating minor nosebleeds can often be managed **at home with simple first aid measures**.
- If needed, the course of treatment can progress to **visiting a physician or the emergency room**.
- In more serious or recurrent cases, **surgical intervention** may be required.

- **Anterior epistaxis** is usually the easiest to manage.
- **Posterior epistaxis** is more likely to require medical attention, since it is more difficult to control and may compromise the airways or cause aspiration.

TREATMENT

AT HOME TREATMENT

- * SIT UP
- * SLIGHTLY LEAN FORWARD and TILT HEAD FORWARD



- * PINCH TIP of NOSE with TWO FINGERS for 15-20 MINS

IF BLEEDING CONTINUES

TOPICAL NASAL SPRAYS



- * VASOCONSTRICTIVE
- * LOCAL ANESTHETICS

IN CASE of:
* RECURRENT EPISODES
* NON-STOP BLEEDING

HOSPITAL SETTING TREATMENT




NASAL PACKING

- * INSERT GAUZE-LIKE MATERIAL or NASAL TAMPON in NASAL CAVITY

IF:
* MEDICATION doesn't WORK
* POSTERIOR EPISTAXIS


TREATMENT

- If we are able to identify the bleeding point: **Cauterization**.
- Can be:
- → Chemical: **Silver nitrate**.
- → Thermal: **Electrocautery device**.




TREATMENT

- If the bleeding site can not be seen then perform: **Nasal Packing.**
- Can be:
 1. **Anterior nasal packing.**
- Involves inserting a gauze-like material or nasal tampon into the nasal cavity to absorb blood and provide pressure to the affected area



TREATMENT

- 2. Posterior nasal packing.
- Can be performed by using:
conventional ribbon gauze or
a Foley catheter.



EPIDEMIOLOGY


- The life long incidence of epistaxis is **60%**, however only **10%** seek medical attention.
- Males > Females.

HISTORY

- Age
- Onset, duration, severity, frequency
- Bilateral or unilateral
- Preceding factors: exercise, sleep, migraine, trauma
- Bleeding from other sites
- Aggravating and relieving factors
- Nasal discharge
- Medical conditions
- Current medications
- Smoking and drinking habits
- Previous epistaxis, recurrent bleeding, easy bruising
- Family history of bleeding disorders

PHYSICAL EXAM


- **General exam:**
examine the skin for evidence of bruises or petechiae that may indicate an underlying hematologic abnormality.
- **Vital signs:**
persistent tachycardia must be recognized as an early indicator of significant blood loss requiring intravenous (IV) fluid replacement and, potentially, transfusion.
- Gently insert a nasal speculum and spread the naris vertically. Start with **inspection**, looking specifically for any obvious bleeding site on the septum that may be amenable to direct pressure or cautery.



LOCALIZATION OF BLEEDING

- Soak some pledgets with **anesthetic-vasoconstrictor solution** and insert them into the nasal cavity to **anesthetize and shrink the nasal mucosa**.
- Allow them to remain for **10–15 minutes**.
- Visualize the cavity with a speculum + a good light source.
- Aspirate excess blood and clots.
- If the bleeding originated from Little's area, it will be clearly visible.
- If an anterior source cannot be visualized, if the hemorrhage is from both nares, or if constant dripping of blood is seen in the posterior pharynx, the bleeding may be from a posterior site.
- Massive epistaxis may be confused with hemoptysis or hematemesis. Blood dripping from the posterior nasopharynx confirms a nasal source.

Approximately 90% of nosebleeds can be visualized in the anterior portion of the nasal cavity.



DIFFERENTIAL DIAGNOSIS

- Allergic rhinitis
- Trauma
- Cocaine toxicity
- Coumarin plant poisoning
- Nasal foreign bodies
- NSAID toxicity
- **Osler-Weber-Rendu syndrome**
- Hemophilia A
- Hemophilia B
- von Willebrand disease
- Warfarin and sub-warfarin toxicity



COMPLICATIONS

- Sinusitis
- **Septal hematoma**/perforation
- External nasal deformity
- Mucosal pressure necrosis
- Vasovagal episode
- Aspiration




PREVENTION

- Correction of bleeding disorders
- Control of hypertension
- Use of humidifiers or vaporizers
- Nasal saline sprays, ointment, Vaseline
- Avoid hard nose blowing or sneezing
- Sneeze with the mouth open
- Avoid nose picking
- Control the use of medications
- Avoid excessive alcohol drinking and smoking



Nasal trauma



CONTENTS

01

Introduction

02

Types

03

Management



04

Complications

INTRODUCTION

- The nose is the most frequently injured facial structure. It may be injured in various forms of sport, in personal assaults and in traffic accidents.
- Nasal deformity depends on **the force and direction of trauma**.
- Injury to the nose may result in one or more of the following:
 - 1- Epistaxis
 - 2- Fractures of the nasal bone
 - 3- Fracture or dislocation of the septum
 - 4- Septal hematoma


INTRODUCTION



TYPES OF FRACTURES


Type 1

only one bone is affected (nasal bone) due to anterior trauma



Type 2

two bones are affected (the frontal process of maxilla and the nasal bone) due to lateral trauma



Type 3

labyrinth of ethmoid
basal skull fracture
base of orbit, maxilla or mandible

MANAGEMENT

1. ABCs
2. Look for **signs** of fracture in the nose (**swelling and discoloration of the skin and subcutaneous tissue covering the nasal bones, tenderness, mobility of the nose, crepitation, obvious deformity**)
3. Look for skull and chest fractures
4. Look for signs of intracranial or abdominal bleeding
5. Admission if the patient is hypotensive, has severe epistaxis,...
6. Imaging:
Plain X-Ray is most commonly done
CT scan if:
 - 1- Type 3 fracture
 - 2- Fracture of other facial bones
 - 3- Evidence of CSF leak




MANAGEMENT

Getting the nose back to its normal shape:

- Presentation **within hours** → the deformity is obvious, you fix the deformity immediately.
- Presentation **after a day** → you can not evaluate the condition due to **edema and hematoma** formation, you wait for one week, but you control the bleeding and give the patient pain killers and antibiotics in case of infection.
- -Presentation **after a month or more** → the process of healing has already started, we wait till it's complete (**6-12 months**), then the patient needs **septorhinoplasty** under GA.

COMPLICATIONS

1. Septal hematoma



2. Septal dislocation



COMPLICATIONS

—septal hematoma


In the setting of trauma to the anterior nasal septum, hematoma formation may occur.

The anterior portion of the nasal septum is composed of a thin cartilaginous plate with a closely adherent perichondrium and mucosa.

A **septal hematoma** is a blood-filled cavity between the cartilage and the supporting perichondrium.

Bacterial proliferation and abscess formation may then result from the presence of stagnant blood. **A hematoma may become infected within 3 days of the trauma.**

Although septal hematomas are uncommon, early diagnosis and treatment is important to **prevent abscess formation, septal perforation, saddle-nose deformity** and potentially permanent complications.



COMPLICATIONS

—septal hematoma


Nasal septal hematoma in:

- **Adults** → typically occurs with significant facial trauma and nasal fracture.
- **Children** → may be found with minor nasal trauma such as simple falls.

COMPLICATIONS

—septal hematoma


- A septal hematoma can usually be diagnosed by **inspecting** the septum with a nasal speculum.
- **Asymmetry of the septum with a bluish or reddish fluctuance** may suggest a hematoma.
- Direct **palpation** may also be necessary, as newly formed hematomas may not be ecchymotic. The best way to palpate is to insert a gloved small finger into the patient's nose and palpate along the entire septum, feeling for swelling, fluctuance, or widening of the septum.
- Most specific image **is CT scan.**



COMPLICATIONS

—septal hematoma

- When a septal hematoma is identified, it should be **aspirated** immediately or incised with the aid of local anaesthesia.





THE END.

Thank you for listening.