

# Introduction to Interventional Radiology

**Dr. Osama Samara.**  
**Consultant Interventional Radiologist**  
**Associate Professor of Radiology**  
**Jordan University Hospital.**



# INTERVENTIONAL RADIOLOGY

A subspecialty in radiology which provides minimally invasive procedures using imaging modalities to diagnose and treat different conditions, it has many different advantages such as.

- Minimally invasive
- Local anesthesia
- Early recovery



# TYPES

1\_ Percutaneous Example biopsy or abcess drainage .

2-endovascular such arterial angioplasty or tumour embolization

# VASCULAR IR

- Endovascular aneurysm repair
- Peripheral arterial angioplasty and stenting
- Thrombectomy and thrombolysis
- Venous thrombolysis
- IVC filter insertion
- Peripherally inserted central catheter
- Embolization
- Transjugular Intrahepatic Portosystemic Shunt
- Uterine fibroid embolization

# PERCUTANEOUS NON-VASCULAR IR

- Abscess drainage
- Percutaneous gastrostomy
- Percutaneous biliary drainage and biliary stent insertion
- Percutaneous nephrostomy
- Image-guided biopsy
- Transcatheter arterial chemoembolization
- Embolization of hypervascular metastases
- Transforaminal nerve root block



# Types Of Vascular Imaging

Doppler  
Ultrasound

Digital  
subtraction

CT  
angiography


MR  
angiography




# TYPES OF ANGIOGRAPHY

- **Advanced US, CTA and MRA techniques made conventional angiography limited to therapeutic purposes.**


# NORMAL ANATOMY









- Arch of aorta
- Subclavian arteries
- Common carotid arteries
- Brachiocephalic trunk






**(b) Anterior view**



## Arterial and Venous Circulation of the Legs



# ANGIOGRAPHY


## Definition

- It is an x-ray examination.
- The blood vessels are opacified by the iodine-containing contrast medium

## Angiograms:


1. Arteriograms
2. ~~VENOGRAMS~~ Venograms





**C. Anteroposterior View**

# Angiogram





# DSA


*(Digital subtraction angiography)*




## ■ Principle

Subtract the shadows that are present on the plain films from the films taken after the contrast has been injected for the angiogram, the result is an image containing details of the opacified structures only, such as arterial system .








Digital vascular imaging  
(2D DSA)



Three dimensional Imaging  
(3DDSA)



# Patient Preparation

- Appointment time
- Nil orally 4-6 hrs.
- On trolley
- In hospital gown
- Groin shave
- Records
- Coagulation profile
- Should be well hydrated.
- Should void before procedure.
- Peripheral pulses marked.
- I.V line in place.
- Informed consent


# EQUIPMENTS




General angiographic room with biplane C-arm digital imaging

# SELDINGER TECHNIQUE


- Site cleaned, area draped, local anesthetic given.
- The **needle** is introduced into the target artery or vein .




Step 1. Insertion of needle  
(with inner cannula)




Step 2. Placement of needle in  
lumen of vessel (inner  
cannula removed)




Step 3. Insertion of guide wire



Step 4. Removal of needle



Step 5. Threading of catheter  
to area of interest



Step 6. Removal of guide wire

**Fig. 21-36.** Six steps of Seldinger technique.

# SELDINGER TECHNIQUE


- A **guide wire** is inserted through the needle.
- With guide wire in vessel, needle is removed.
- **Catheter** is threaded onto the guide wire.
- Under fluoroscopy, the catheter is then advanced and the guide wire is removed.





# Femoral Artery Approach

- Easily accessible.
- Large caliber vessel
- Well defined land marks exist.
- Low complication rate.



# Post procedural Care

- After the catheter is removed **compression** is applied to the puncture site.
- The patient is asked for **bed rest** for a minimum of 4 hours
- During rest patient is monitored **and vital signs** like peripheral pulses distal to Puncture are regularly checked
- The extremity is also checked for warmth, color, numbness to **ensure circulation** has not been disrupted.

# CONTRA-INDICATIONS

## ■ Contra-indications to contrast media

1. **Allergy (allergy to CM is an absolute contraindication)**

2. **Impaired renal function**

3. **Medications**



■ **Blood- clotting disorders/ Anti coagulant medication**

■ **Unstable cardio pulmonary/ neurological status**

# COMPLICATIONS

■ **0.16% major complication rate:**

- 1. Local complications:** hematoma, vessel laceration, dissection, pseudoaneurysm, AV fistula.
- 2. Systemic complications:** contrast reactions, fever, sepsis, dehydration, death.
- 3. Therapy related complications** (ex: **CNS bleeding during thrombolysis**)



## ■ **Peripheral Vascular Disease**

■ **Aneurysms**

■ **Stenosis**

# Arterial Occlusive Disease

- Acute VS chronic.

# Acute Arterial Occlusion

- **Embolus** is the most common cause
- Other etiologies are dissections of the arterial wall, external compression , spasm or hemodynamic problems.
- **Immediate intervention** needed to prevent loss of extremity




# Acute Embolic Occlusion

- Cardiac origin : recent MI or A-fib
- Emboli get trapped at **arterial bifurcations** and cause further thrombosis by apposition or stagnation of blood flow
- Most frequent location of arterial emboli is **the common femoral artery (46 %)**

# Acute Thrombotic Occlusions

- Caused by atherosclerosis in over 90%.
- The clinical signs are less severe than with acute embolic disease.

Normal artery



Blood flow

Artery narrowed by atherosclerosis



Plaque



```
graph LR; PVD[PVD] --- Occlusion["Occlusion (Thrombotic/Embolic):  
Felling defect, meniscus sign,  
Sharpe cut off or Non  
visualization"]; Occlusion --- Acute[Acute]; Occlusion --- Chronic[Chronic]
```



**PVD**

**Occlusion  
(Thrombotic/Embolic):**

Felling defect, meniscus sign,  
Sharpe cut off or Non  
visualization

Acute


Chronic



**Acute occlusion**

# Acute Occlusions

- On arteriography an acute thrombotic occlusion may show a relative **sharp cut-off** of the contrast column



LT



ROT  
-1

ANG  
-1

RUN  
2  
20  
MASK IMAGE  
2 12




Aortoiliac Occlusive Disease, Also Known As  
Leriche's Syndrome



Angioplasty and stenting of superficial femoral artery

# ANEURYSMS






# ANEURYSMS

## Dilation of an artery:

1. Fusiform or saccular
2. True or false


## Aorta :

- Thorax > 4cm.
- Abdomen > 3cm.



# TRUE ANEURYSMS

- localized outpouchings, spindle shaped which involve **all three layers** of the arterial wall.
- The pathogenesis is mainly a degeneration of the media
- The etiology is atherosclerosis in 70% to 80%.



# ANEURYSMS



# RUPTURED ANEURYSM



# Dissecting Aneurysms



# Arterial Stenosis

Artery stenosis may be caused by several pathological processes:

- Atherosclerosis (~75% of cases).





# VENOGRAPHY

## ■ Technique

contrast medium is injected into a small vein in the dorsum of the hand or foot.

**LIMITED**







# DVT

```
graph TD; DVT[DVT] --- Central[1. Filling defect<br/>2. Non-visualization]; Central --- Acute[Acute]; Central --- Chronic[Chronic:<br/>1. Collaterals<br/>2. Faint opacification];
```

The diagram is a flowchart with a central box containing two items: '1. Filling defect' and '2. Non-visualization'. A box labeled 'DVT' is connected to the top of this central box. Two other boxes are connected to the sides of the central box: 'Acute' on the left and 'Chronic: 1. Collaterals, 2. Faint opacification' on the right. The top of the page features a horizontal bar with three segments: blue, orange, and yellow.

1. Filling defect

2. Non-visualization




Acute

**Chronic:**

1. Collaterals

2. Faint opacification






A

B

C



# Interventional Radiology


- **The interventional radiologist uses **per-cutaneous techniques under imaging control** to guide small instruments through the blood vessels or other pathways to treat disease.**
- **These procedures are typically much less invasive and cause much less discomfort.**
- **Imaging modalities:**
  1. **X-ray /fluoroscopy/DSA**
  2. **CT**
  3. **US**
  4. **MRI**

# Basic Interventional Techniques

1. Endovascular Aneurysm Repair
2. Angioplasty
3. Vascular Stents And Filters
4. Therapeutic Embolization
5. Percutaneous Needle Biopsy
6. Percutaneous Abscess drainage or aspiration

# I. Endovascular Aneurysm Repair

- For high risk surgery patients
- Before aneurysm reaches diameter for elective surgery
- Inserted through femoral artery
- Decreased length of stay in hospital
- Still need monitoring for complications



## 2. Angioplasty

### ■ Technique

During the procedure a balloon catheter is placed across the lesion and the balloon is inflated and the lesion is compressed. The result is a wider vessel lumen and increased blood flow.

### ■ Indications

Short vascular stenosis or occlusion in the legs or arms, kidneys, brain or elsewhere in the body.






Diagram of angioplasty





Angioplasty of left femoral artery

# 3. Vascular Stents

## ■ Technique

Stents are expandable metal cylinders that can be embedded in plastic and collapsed to enable them to be inserted through an artery or vein. Stents help hold the artery open, which improves blood flow and relieves symptoms caused by the blockage.

## ■ Indications

Vascular stenosis in peripheral vascular





Diagram of stenting

## Indications For Stents In Revascularization Procedures:

- Unsuccessful PTA (percutaneous transluminal angioplasty)
- Recurrent stenosis after angioplasty.
- Long segment stenosis.
- Total occlusion.
- Hard calcified or ulcerated plaque.
- Renal osteal lesions.



**Angioplasty of left subclavian artery**




I V C Filters

# Inferior Vena Caval Filters


- Inferior vena caval filters can be introduced percutaneously through the femoral vein.
- The filters trap emboli originating from leg or pelvic vein thrombi.
- They are used in patients who are at risk of pulmonary embolism.

Site of Pulmonary Embolus



© 2003 Society of Interventional Radiology

Deep Vein Thrombosis (DVT)



© 2003 Society of Interventional Radiology

# Indications For IVC Filters:

DVT and/or PE and one of the following:

1. Contraindication to anticoagulation.
2. Failure of anticoagulation.
3. Complication of anticoagulation.

# Technique:


- Through right femoral vein.
- Because caval thrombosis is a complication of filter placement, filters are usually placed below the renal veins.



# Complications:

- Filter migration.
- Filter failure (recurrent PE).
- IVC thrombosis
- Groin complications.
- PE after IVC filter may be due to filter thrombosis, collaterals, upper extremity DVT.

# Inferior Vena Caval Filters




# 4. Therapeutic Embolization

## ■ Technique

Introducing a variety of materials through a catheter selectively placed in the vessel.




## ■ Indications

1. Bleeding
2. Tumors
3. Aneurysms
4. Arteriovenous malformations




## Gastrointestinal (GI) tract bleeding

Angiography demonstrated the bleeding site at left gastric artery.



## Hemoptysis


1. The extravasation of contrast medium, local hypervascularity and shunting of contrast from bronchial arteries to pulmonary artery are demonstrated in angiography of bronchial arteries
2. Following embolization with gelatin foam, the bronchial artery is occluded.



*(A) Right frontal meningioma (tumor); (B) Arteriogram showing blood supply to tumor prior to embolization; (C) Arteriogram after embolization of blood supply to tumor.*

## **Preoperative tumor embolization**

- Many tumors such as meningiomas and renal cell metastases are hypervascular thus making surgical resection more difficult and time consuming.
- Embolization of the tumor preoperatively with gelatin foam, particles, coils, and alcohol can make resection easier and faster.







# Transarterial chemoembolization

## Technique

- Chemotherapeutic agents are usually combined with embolic particles to achieve chemoembolization of hepatic tumors.
- The aim is to cause ischemia and prolonged contact of the chemotherapeutic agent with the tumor.
- Such mixtures can dramatically increase the local concentration of the chemotherapeutic agent.
- The systemic drug levels is lower, thus reducing toxicity.

## Survival

- The probability of cancer recurrence and/or metastatic dissemination was lower after TACE than after surgery.



## Cerebral aneurysms

Carotid angiogram shows an aneurysm located in the Acom A.

The embolization of aneurysm with metal coils is well demonstrated.



## Complications Of Embolization:

- Post-embolisation syndrome (fever, elevated WBC).
- Infection of embolised area (prophylactic antibiotics).
- Reflux of embolic material (non targeted embolization).
- Alcohol can cause skin, nerve and muscle necrosis .

# 5. Percutaneous Needle Biopsy

## Technique

- A needle is passed into the desired site and a small amount of tissue is removed.
- With a fine needle(20-22 gauge for aspiration, 14-18 gauge for soft tissues,
- 10-13 gauge for bone.

## Indications

- Diagnostic test for breast, lung and other cancers
- An alternative to surgical biopsy.




## 6. Percutaneous Drainage Of Abscesses And Other Fluid Collections




### Technique

- Specially designed drainage catheters can be introduced percutaneously into abscesses allowing the pus to drain for several days.

### Indications

- Intra abdominal abscesses





Percutaneous Insertion Of biliary Stent To Bypass An Obstruction In The biliary Ducts



Thank You !