

MULTIPLE PREGNANCY

Dr Amal Barakat

JU/5th Year Medical Students

October/2023

MULTIPLE PREGNANCY

- A pregnancy with more than one fetus is called a multiple pregnancy

TERMINOLOGY

- Two offspring – [twins](#) //Twins are by far the most common form of multiple births in humans
- Terms used for the number of offspring in a multiple birth, where a number higher than three ends with the suffix *-uplet*:
 - three offspring – triplets
 - four offspring – quadruplets
 - five offspring – quintuplets
 - six offspring – sextuplets
 - seven offspring – septuplets
 - eight offspring – octuplets
 - nine offspring – nonuplets
 - ten offspring – [decuplets](#)

Incidence and epidemiology

- Twin pregnancies and higher-order multiple births comprise an increasing proportion of the total pregnancies in the developed world.
- The incidence of multiple gestation **continues to increase**, and now accounting for more than **3%** of all live births.

Incidence and epidemiology

- The U.S. [Centers for Disease Control and Prevention](#) report more than 132,000 sets of twins out of 3.9 million births of all kinds each year, about 3.4%, or 1 in 30.
- Without fertility treatments, the probability is about 1 in 60; with fertility treatments, it can be as high as 20-25%.
- **DZ twins are more common than MZ twins, 69 and 31 % of twins, respectively.**
- **The incidence of MZ twins is relatively stable worldwide at 3 - 5 /1000 births**

Rising multiple birth rate is due mainly to:

- **Expanding use of Fertility drugs** (Drugs used for induction of ovulation)
- **Increasing use of assisted reproduction techniques, including in vitro fertilisation (IVF).**
- *Up to 24% of successful IVF procedures result in multiple pregnancies*
- **Increasing maternal age at conception.** Older women are more likely to become pregnant with multiples.
- **Changes in population demographics** (due to immigration) have also contributed to the rise.

MATERNAL COMPLICATIONS

- Multiple gestations are associated with significantly higher risks for:
 - ✓ Hypertension
 - ✓ Placental abruption
 - ✓ Preterm labor
 - ✓ Preeclampsia (26%);
 - ✓ HELLP syndrome (9%)
 - ✓ Anemia (24%)
 - ✓ Preterm premature rupture of membranes (preterm PROM) (24%)
 - ✓ Gestational diabetes (14%)
 - ✓ Acute fatty liver (4%)
 - ✓ Chorioendometritis (16%)
 - ✓ Postpartum hemorrhage (9%)
 - ✓ Miscarriage
 - ✓ Operative Delivery
 - ✓ Postnatal illness

MATERNAL COMPLICATIONS

- The risk of pre-eclampsia for women with twin pregnancies is almost three times that for singleton pregnancies, while the risk for triplet pregnancies is increased nine-fold.
- In addition, maternal mortality associated with multiple births is 2.5 times that for singleton births.

FETAL COMPLICATIONS

- Prematurity
- Monochorionicity with its complications
- Growth restriction.
- ^^Stillbirth
- Infant mortality rates increase
- long- term morbidity increase (especially neurodevelopmental disability and chronic lung disease)
- The incidence of severe handicap among neonatal survivors of multiple gestation is also increased.
- Major congenital abnormalities are 4.9% more common in multiple pregnancies than in singleton pregnancies.

PRETERM & TWIN PREGNANCY

- **Preterm birth** occurs in more than **50% of twin** and **75% of triplet** gestations.
- Duration of pregnancy becomes shorter with increasing numbers of fetuses.
- The mean duration of pregnancy is **35.3 weeks for twin gestations, 31.9 weeks for triplets, and 29.5 weeks for quadruplets.**
- The higher incidence of preterm birth in multiple pregnancies is associated with an increased risk of neonatal mortality and long- term morbidity (especially neurodevelopmental disability and chronic lung disease).
- A cervical length of less than **20 mm** in a **twin pregnancy** at **20 to 24 weeks** gestation was associated with a **10-fold** positive likelihood ratio for preterm birth before **32 weeks** gestation.
- **Cervicovaginal fetal fibronectin** assay can be used to predict preterm labor.

PRETERM & TWIN PREGNANCY

- **Interventions to prevent preterm labor and prolong pregnancy** for patients with multiple gestations :
 - ✓ prophylactic **cervical cerclage** (????)
 - ✓ Supplemental **progesterone** (????)
 - ✓ **Bedrest** (????)
 - ✓ **Tocolytic drugs** (????)

Zygoty

- Dizygotic (DZ), resulting from the fertilization of two separate ova during a single ovulatory cycle.
- DZ twins have dichorionic-diamniotic (DCDA) placentas.
- Dizygotic twins are always diamniotic, dichorionic (i.e., have 2 sacs and 2 placentas). The 2 placentas may fuse but do not have vascular connections.

Factors influencing the incidence of DZ twins are:

- **Factors influencing the incidence of DZ twins are:**
 - ✓ **Use of fertility stimulating drugs**(twin births increased from 1/53 infants in 1980 to 1/30 infants in 2009)
 - ✓ **Maternal age**(One-third of the increase in multiple births in recent decades has been attributed to increasing age at childbirth)
 - ✓ **Race/geographic area** (1.3/1000 Japan, 8/1000 United States and Europe, 50/1000 Nigeria)
 - ✓ **Parity**
 - ✓ **Family history**
 - ✓ **High BMI & Maternal height**


Zygoty

- Monozygoty (MZ), resulting from a single fertilized ovum that subsequently divides into two separate individuals.
- Monozygoty (identical) twins do not run in families.
- Monozygoty twins have different amnionity and chorionity depending on the stage of cleavage of the single fertilized ovum.


Zygoty

- **Monozygotic twins have different amnionity and chorionity depending on the stage of cleavage of the single fertilized ovum**
- **In MZ Twin the timing of egg division determines placentation**
 - **Diamniotic, dichorionic (DCDA) placentation occurs with division prior to the morula stage (within 3 days post fertilization). This occurs in 29% of the cases.**
 - **Diamniotic, monochorionic (MCDA) placentation occurs with division between 4-8 days postfertilization. This occurs in 70% of the cases.**
 - **Monoamniotic, monochorionic (MCMA) placentation occurs with division between 8-12 days postfertilization. This occurs in 1% of the cases.**
 - **Division at or after day 13 results in conjoined twins. This is an extremely rare condition occurring in up to 200,000 births**

In MZ twins, the timing of egg division determines placentation



Division at or after day 13 results in conjoined twins.



Conjoined Twin

- A subset of **monozygotic twin gestations** in which **incomplete embryonic division** occurs **13 to 15 days** after conception, resulting in varying degrees of fusion of the two fetuses.
- Classified according to the **anatomical site of union** (eg, chest, head).
- **Associated congenital defects** unrelated to the area of fusion are common, as is stillbirth.
- **Delivery of viable infants is always by cesarean.**

Ultrasound in Twin pregnancy

- **Diagnosis:**

- **Clearly separate gestational sacs, each surrounded by a thick echogenic ring, is suggestive of dichorionicity.**
- **Visualization of multiple gestational sacs with yolk sacs by 5 weeks, or multiple embryos with cardiac activity by 6 weeks.**

Clearly separate gestational sacs, each surrounded by a thick echogenic ring, is suggestive of dichorionicity.



Visualization of multiple gestational sacs with yolk sacs by 5 weeks, or multiple embryos with cardiac activity by 6 weeks.



Ultrasound in Twin pregnancy

- All women with a twin pregnancy should be offered an ultrasound examination at **10–13 weeks** of gestation to assess: **viability, chorionicity, major congenital malformation** and **nuchal translucency**.
- **Careful sonographic surveys of fetal anatomy** are indicated in multifetal pregnancies, because the risk for **congenital anomalies** is **increased (3-5 fold higher)**.

Ultrasound in Twin pregnancy

- **After confirming a diagnosis of multiple gestation we should determine chorionicity.**
- **The accuracy of ultrasound in the assessment of chorionicity in the first trimester is high and is mainly based on the evaluation of lambda and T-sign**

Ultrasound in Twin pregnancy

- **Fetal growth should be assessed by serial ultrasonography.**
- **Intrauterine growth of twins is similar to that of singletons until 30 - 32 weeks gestation.**
- **Assess for growth discordance (discordance greater than 20% has also been shown to be an important predictor for adverse perinatal outcomes)**
- **Assess cervical length in multiple gestation to identify those at increased risk for preterm delivery.(16 -24week , A cut-off of 20 mm)**
- **A cervical length of less than 20 mm in a twin pregnancy at 20 to 24 weeks gestation was associated with a 10-fold positive likelihood ratio for preterm birth before 32 weeks gestation.**

Chorionicity

- **Chorionicity is the main determinant of the perinatal outcome in twin pregnancies:** perinatal mortality and morbidity are significantly higher in monochorionic versus dichorionic twins.
- **This is mainly due to complications associated specifically with monochorionicity such as:**
 - **Twin to Twin transfusion syndrome (TTTS),**
 - **Selective Fetal Growth Restriction (sFGR)**
 - **Twin Reverse Arterial Perfusion Syndrome (TRAP)**

Chorionicity

- The prenatal determination of chorionicity is the first step for an accurate managing of twin gestation.
- Knowledge of chorionicity helps in **risk assessment, genetic counseling, invasive procedure and management of TTTS and selective IUGR, death of one twin and discordant fetal anomaly.**
- It is best done in the first trimester, when the diagnostic accuracy approaches 100%.
- The most reliable sonographic signs are the lambda and T-sign and the number of the placental masses evaluated before 14 weeks of gestation

Chorionicity

- **If it is difficult to determine chorionicity even after referral:**
- *Manage the pregnancy as monochorionic until proved otherwise.*

Fetal gender

- **The identification of discordant fetal gender indicates dichorionic twinning.**
- **The positive predictive value of discordant gender (when correctly identified) is 100%.**
- **However, considering that around 50% of concordant sex twins are dichorionic, the definition of chorionicity needs further sonographic signs.**

Different sonographic signs may be used to evaluate chorionicity:

- **Different sonographic signs may be used to evaluate chorionicity:**
- 1- number of placental masses
- 2- sex of the fetuses. The identification of discordant fetal gender indicates dichorionic twinning
- 3- Membrane thickness: **membrane thickness of 2 mm** helps in diagnosing **chorionicity**.
- 4- Number of membrane layers: **placentation is MC** if only **two layers** are present; the presence of **three or four layers suggests dichorionicity**.
- 5- characteristics of the intertwin membrane is the most useful and valuable tool: the take-off of the membrane from the placental surface shows the typical “lambda” appearance in dichorionic pregnancy and the typical “T” appearance in the monochorionic ones.

The twin peak sign (the lambda (λ) sign)

- **Seen in Dichorionic Diamniotic twin pregnancy.**
- **This sign is a triangular projection of placental tissue which extends from the placenta between the layers of amniotic and chorionic membranes of each fetus**
- **It is best seen in the first trimester (between 10-14 weeks)**

The twin peak sign (the lambda (λ) sign)



Diagn Prenat. 2014;25:58-64

The T sign

- Seen in Monochorionic Twin
- It refers to the lack of chorion extending between the layers of the intertwin membrane, and the appearance of the **thin intertwin membrane** as it takes-off from the placenta at a **90 degree angle**, denoting a monochorionic pregnancy.

The T sign



Diagn Prenat. 2014;25:58-64

ANTEPARTUM CARE AND FETAL SURVEILLANCE

- **Serial sonographic assessment of fetal growth** is recommended in multiple gestations.
 - ✓ every **3 - 4 weeks** from **18 weeks gestation** in **DC twins**, or every **2 weeks** if growth restriction or growth discordance ($>20\%$) is discovered.
 - ✓ **MC twins**, as well as all **higher-order multiple gestations**, serial growth scans are performed every **2 weeks** from **16 weeks** gestation.
- When **significant growth discordance** is confirmed, fetal testing should begin intensively. This consists of **twice-weekly NST** supplemented by **biophysical profiles** and **umbilical artery Doppler velocimetry**.

Antenatal corticosteroid prophylaxis for reducing perinatal morbidity

- **A single course of corticosteroids is recommended for pregnant women between 24 0/7 weeks and 33 6/7 weeks of gestation who are at risk of preterm delivery within 7 days, including for those with ruptured membranes and multiple gestations.**

Timing and Method of delivery

- The Royal College of Obstetricians and Gynaecologists (RCOG) and National Institute for Health and Care Excellence (NICE) recommend uncomplicated MCDA twins to be delivered between **36** and **37** weeks and DCDA twins to be delivered between **37** and **38** weeks.
- In general, All **twin fetuses** should be **delivered by 39 weeks** of **gestation** because of the **rising perinatal morbidity and mortality** beyond that date.
- In **MCMA twins** delivery at about 32-34 weeks **weeks should be considered** because of the **increasing risk of perinatal mortality and unexpected fetal loss in the third trimester**.

Timing and Method of delivery

- **Fetal presentations** and size of twin should be determined before choosing the mode of delivery.
- **Electronic fetal heart monitoring** should be available for both.
- **Epidural anesthesia is recommended**

Timing and Method of delivery

- For **vertex-vertex presentation** and in the **absence of obstetric indications for cesarean delivery**, **vaginal birth** should be planned regardless of gestational age.
- There is **no absolute indication** to deliver the **second twin** within a **specified time limit (continuous FHM)**.
- For **vertex-non-vertex twins**, **Vaginal delivery allowed with breech delivery of the 2nd twin**.
- If the 2nd twin is significantly **larger than the 1st**, **cesarean delivery** is recommended.
- For **non-vertex 1st twin** **cesarean delivery** is recommended.

Timing and Method of delivery

- **Higher-Order Multiple Gestations:**
 - ✓ **cesarean delivery** under **regional anesthesia** for all patients with three or more live fetuses that are of a viable gestational age is recommended.
- **Monoamniotic twins:**
 - ✓ **Cesarean birth** is recommended to avoid complications from **cord entanglement**.

Cesarean section or vaginal delivery??

- **In Summry:**
- **Cesarean delivery in multiple pregnancy is needed when:**
 - First twin is in non cephalic presentation
 - When it is a Monoamniotic twin pregnancy.
 - Higher-Order Multiple Gestations (Triplets, ..)
 - If there is medical or obstetric indication

Feto-fetal transfusion syndrome (FFTS)

- FFTS occurs in **15% of MC twin** pregnancies, and accounts for about 20% of stillbirths in multiple pregnancies.
- TTTS occurs almost exclusively in monozygotic twins with monochorionic-diamniotic (MCDA) placentation.
- It can develop at any point during pregnancy but typically emerges in the second trimester.
- Vascular communication is present virtually in all monochorionic twins; in most cases, the blood flow is balanced with no net transfusion of blood from one twin to the other
- **TTTS** occurs because of an **imbalance in blood flow** through **vascular communications in the placenta**, which leads to **overperfusion** of one twin and **underperfusion of its co-twin**.
- **Arterio-venous unidirectional anastomoses** result in net transfusion of blood from the donor to the recipient fetus.
- Hypovolemia causes a decrease in the urine output leading to oligohydramnios or anhydramnios, while hypervolemia in the recipient twin causes an increase in the urine output that leads to polyhydramnios.


Feto-fetal transfusion syndrome (FFTS)

- **Ultrasonographic criteria for diagnosis of TTTS include :**
 - ✓ Presence of a single placenta
 - ✓ Sex concordance
 - ✓ Significant growth discordance (approximately 20%)
 - ✓ Discrepancy in amniotic fluid volume between the two amniotic sacs (usually oligohydramnios and polyhydramnios)
 - ✓ Presence of fetal hydrops or cardiac dysfunction
 - ✓ Abnormal umbilical artery Doppler findings, such as absent end-diastolic flow in the donor fetus

Feto-fetal transfusion syndrome (FFTS)


- **The donor fetus** is **hypoperfused**, demonstrating signs of **intrauterine growth restriction**, **anaemic** and **oligohydramnios**.
- **The recipient fetus** is **hyperperfused**, **hypertensive**, demonstrate **biventricular hypertrophy and diastolic dysfunction**, and **polyhydramnios**.
- FFTS is associated with a **high risk of fetal/neonatal mortality**, and fetuses who survive are at risk of **severe cardiac, neurologic, and developmental disorders**.
- management approaches for the treatment of **severe TTTS** (24 to 26 weeks) gestation:
 - ✓ **Serial reduction amniocenteses**
 - ✓ **Amniotic septostomy**
 - ✓ **Selective fetoscopic laser coagulation of placental anastomoses.**
 - ✓ **Fetoscopic Laser Photocoagulation (FLP)** is the best treatment option for stages 2 through 4 and is approved by the Food and Drug Administration (FDA) for use during 16 to 26 weeks' gestation.

Feto-fetal transfusion syndrome (FFTS)



Twins with twin-twin transfusion syndrome

Feto-fetal transfusion syndrome (FFTS)




Fetal laser photocoagulation

Twin Reversed Arterial Perfusion Sequence(TRAP)

- Occur in **1%** of **MZ twins**.
- One twin has an **absent, rudimentary, or nonfunctioning heart (acardiac)**.
- **The donor (pump twin)** provides circulation for itself and for the **recipient (perfused twin)** through a **direct arterio-arterial anastomosis** at the placental surface.
- The pump twin is at risk for development of **hydrops** or congestive **cardiac failure**.

Twin Reversed Arterial Perfusion Sequence(TRAP)

Twin Reversed Arterial Perfusion (TRAP) Sequence



Intrauterine Demise of One Fetus

- Loss of one twin in the first trimester does not appear to impair the development of the surviving twin.
- Fetal death occurring after mid gestation (17 weeks) may increase the risk of intrauterine growth restriction (IUGR), preterm labor, preeclampsia, and perinatal mortality.
- After the death of one twin in a MC gestation, approximately 15% of remaining fetuses also die, compared with approximately 3% of remaining fetuses in a DC gestation.
- The risk for significant neurologic morbidity is increased after intrauterine death of one fetus in a MC, but not in a DC gestation.
- Maternal coagulopathy, the most feared complication following twin demise, appears to be uncommon. However, coagulopathy has been reported to occur about 3–5 weeks following fetal demise.

Multifetal Pregnancy Reduction(MFPR)

- The incidence of **high order multifetal gestation** (ie, triplets or more) increased dramatically, due to widespread use of assisted reproductive technology (ART).
- These pregnancies are at higher risk of **maternal, fetal, and neonatal complications than singleton pregnancies**
- Higher order multifetal gestations **should be prevented** by better control of ovulation induction and embryo transfer.
- It is recommended to use single embryo transfer in all situations if a top-quality blastocyst is available
- Selective reduction was developed in the mid-1980s, as people in the field of assisted reproductive technology became aware of the risks that multiple pregnancies carried for the mother and for the fetuses.

Multifetal Pregnancy Reduction

- **MFPR** is defined as a first-trimester or early second-trimester procedure for reducing the total number of fetuses in a multifetal pregnancy by one or more.
- In most cases, the involved gestations will be higher-order multifetal pregnancies, defined by the presence of three or more fetuses.
- Operative techniques that may be used are chemical, thermal, radiofrequency and laser, depending on chorionicity as well as other factors.
- Intracardiac potassium chloride is appropriate to employ when there is independent chorionicity. It carries a lower risk of pregnancy loss.
- Vascular occlusion using radiofrequency ablation, bipolar coagulation or intrafetal laser can be employed in monochorionic fetuses and twin reversed arterial perfusion pregnancies, but carry a higher risk of pregnancy loss.

Multifetal Pregnancy Reduction(MFPR)

- **MFPR** is usually performed between **10 and 13 weeks** of gestation.
- Under **continuous ultrasound guidance**, a **needle** is placed into the thorax of the targeted fetus, **2 to 3 mL of potassium chloride** is injected, and **asystole** is observed for at least **3 minutes**.
- Potassium chloride injection must not be used for MFPR in a **single fetus of a MC pair** because of the risk for co-fetal demise or neurologic injury.
- A lot of medical, ethical and religious issues where raised against this procedures.