

## بِسَمِ ٱللهِ ٱلرَّحْمَنِ ٱلرَّحِيمِ

1-What makes the patient avoid anything that stimulate his memory?  Avoidance behavior
2-Patient who experience intense fear when the door is slammed very hard?  Hyper vigilance
3-Young women with a presentation of recurrent palpitation, sweating, shortness of breath and fear of going to die for 3 months, happens in different situation and results that the patient fear leaving her safe place, what is your diagnosis?  Agoraphobia
4-22 years old male came home from night out with his friends acting weird, his family brought him to ER because he fought with his brother saying he was the Devil and wants to kill him, the psychopathology in this case? Delusions.
5-Patient refuse to take oral anti-psychotics, what would you do?  Give IM injection
6-A 70-year-old male patient, lost his spouse started to feel very sad for the last 2 months, had some disorganized behaviors, started to believe he was responsible for the failure of the Arab Spring Protest, what is the psychopathology?  Delusions of guilt
7-Most common hallucination in schizophrenia?  Auditory
8-40 years old patient brought to the clinic by his family. The patient is sure that his wife is cheating on him without evidence, the psychopathology in this case?  Delusion of jealousy or delusion of infidelity
9-Abrupt Withdrawal of one of the following drugs causes seizures: (apparently benzodiazepine withdrawal) ** *Answer: Lorazepam

 ${\bf 10\text{-}obsessive-compulsive\ disorder\ patients\ Have\ the\ following\ characteristics\ Except:}$ 

a)Intrusive
b)Ego-Dystonic
c)Resisted
D) Delusional
Answer: Delusional (not a feature of OCD)
11-A schizophrenic patient want to be pregnant, what is the choice that her off-spring will have Schizophrenia: ***
Answer: 12%
12-One of the following is Wrong about Anxiety disorder:
A. Behavioral therapy used in anxiety disorder.
B. more common in higher socioeconomic groups
c)Anxiety prevalence rate is 10%
Answer: B
13-One of the following is true about Impulse control disorder:
A. Kleptomania is a criminal behavior.
B. Adults are more committed to Pyromania.
C. Serotonin dysregulation may be a cause.
D. Trichotillomania is more common in males.
Answer: C
14-Which of the following is true about adjustment disorder: **
A. a rare mental disorder
B. Usually following a catastrophic Traumatic events
C)Onset of symptoms within 3 months following trauma exposure.
D)Rarely seen in children or adolescent.
E) Conduct abnormalities are infrequent presenting feature in children
Answer is c.
15-what is true about Bipolar disease:
Answer: lithium is the main long-term treatment
46 About ashironhamia
16-About schizophrenia:
Answer: Hospitalization indicated when the patient refuse to eat or drink
17-Which of the following unlikely to be a diagnostic feature of dementia**

- A. Impaired memory
- B. Fluctuating level of Consciousness
- C. Impaired digit span
- D. Difficulty dressing
- E. Difficulty in finding home.

Answer: B) Fluctuating level of consciousness (not a diagnostic feature of dementia)

18-All of the following are true about suicide, Except:

- A) Depression is associated with high rates of suicides.
- B) Suicide is more common among men than women.
- C)Suicide is associated with living alone.
- D) Mild Dementia is a risk factor for suicide
- E) The elderly is less successful in his or her attempt at suicide

Answer: E) The elderly is less successful in his or her attempt at suicide (false about suicide)

19-one of the following is a common side effect of SSRI's: \*\*

- A) Hypotension
- B) Dystonia
- C & D other SE of Antipsychotics
- E) Nausea

Answer: E) Nausea (Common side effect of SSRI)

20-One of the following is a Paraphilia:

Answer: Voyerurism

21-A 23-year-old male consulted his doctor about recurrent short episodes of depression. The doctor prescribed Fluoxetine. After 3 weeks of regular daily use of the drug the patient started to display overactivity, aggressiveness, Overtalkativeness, overspending of money and started to take alcohol in excess. The most likely underlying reason for the recent development was that the doctor missed to make the diagnosis of:

- A) Schizophrenia
- B) Antisocial personality disorder
- C) Alcohol abuse
- D)bipolar disorder
- E) Delirium

Answer: Bipolar disorder

22-A 30-year-old male arrested several times during the last year by police because of driving offences. Screening the blood revealed alcohol in large concentration but no other substances. He gave a history of infrequent heavy drinking during weekends followed by problems with family and neighbors. After 2 weeks of quitting alcohol, he displayed no withdrawal symptoms. The most appropriate diagnosis is:

- A) Alcohol intoxication
- B) Alcohol dependance
- C) Alcohol tolerance
- D)Alcohol abuse
- E) Alcohol Blackout

**Answer: Alcohol abuse** 

23-One of the following drugs is Antipsychotic:

Quetiapine

Rest of choices are Antidepressants.

Answer: Queriapine

24-A 65-year-old woman who has been a known patient of depression for years is brought to emergency room in a state of confusion. She was recently prescribed diazepam by her physician. On examination she is confused, distracted and has a trouble conversing. She is agitated and is disoriented in space and time. According to the attendant, she was fine a few hours ago and has never suffered a similar episode. which of the following is the most likely diagnosis:

- A) Dementia
- B) Delirium
- C)Schizophrenia
- D)Epilepsy
- E) Bipolar disorder

Answer: Delirium

25-one of the following is true: \*\*\*

- a) Sexual identity is the self-identification of gender.
- b) Gender Identity is the Biologic identification of gender.

C)Gender identity usually develops by age 3 (36 months)

Answer: Gender Identity develops by age 3

26-what is the most appropriate statement to describe Delusion:

- a) False belief amenable to reasoning
- b) Thought content disorder.
- c) Misinterpretation of external stimuli

Answer: B.Thought content disorder

27-All of the following is true except:

Answer: stigmatization for psychiatric illnesses occur in developing countries only

Answer: Feeling of unreal self
29-All of the following occurs in Narcolepsy except:  a) Cataplexy b) Sleep paralysis c) Partial seizures  Answer: C) Partial seizures (Does not occur in narcolepsy)
30-one of the following is cluster A personality disorders: Schizoid
31-Wrong about dementia: fluctuations of concentration
32-Wrong about insomnia: 2 episodes per month for diagnosis
33-Most potent on D2: Haloperidol
<ul> <li>34-One of the following regarding suicide is wrong:</li> <li>a) It's the 3rd leading cause of death.</li> <li>b) Mild dementia is a risk factor.</li> </ul> Answer: A
35-Indication for admission bipolar patient: not eat or drink
36-All of the following is true except: stigmatization for psychiatric illnesses occur in developing countries only.
37-One of the following drugs cause seizures if withdrawn: Diazepam
<ul><li>38- When giving lithium for treatment you would do what test:</li><li>a) Serum cortisol</li><li>b) Serum creatinine</li></ul>

28-Depersonalization is:


a) Dementia

Answer) Serum creatinine \*\* (follow up when taking lithium)

39- Social phobia is one of the following: a) Fearing of strangers in open places b) Fearing of negative evaluation \*\* Answer: B 40- indication for hospitalization (in bipolar): a pt. who is not eating or drinking. 41- Wrong about ECT: it's always a 2nd line of treatment. 42- Not given to an elderly with dementia: **TCA** 43- What's wrong: kleptomania is criminal behavior. 44- Risk of schizophrenia in fetus with a pregnant woman? Answer :12% 45- What is depersonalization: Feeling of unreal self 46- What is affect: elicited by the doctor. Objective assessment 47- Not associated with narcolepsy? Night seizures 48- A pts came to the doctor with depression, he was given an antidepressant then a few weeks later he came again with euphoria, agitation (can't remember exactly but they are symptoms of mania) what's the diagnosis? **Bipolar** 49- A 65-year-old women appeared to be confused at the ER. She has been prescribed benzodiazepines. This is her first episode what's this.

b) Delirium	
Answ	er: B
50- A manifestation on Neuroleptic Malignant Syndrome:	
Leukocytosis	
51- A patient is known to drink alcohol and have had high levels in his blood. His drinking led to family problem. When they cut his drinking, he had no withdrawal symptoms what this is called:	ns.
a) Alcohol dependence	
b) Alcohol abuse	
c)Alcohol intoxication	
Answ	er: B
52- Delusion:	
false fixed thought	
53- Social phobia:	
fear of negative evaluation	
54- About schizophrenia:	
it common in winter	
55- What drug causes dryness of mouth:	
Imipramine (TCA)	
56- For lifetime treatment of bipolar depression we use:	
a) Valproate **	
b) TCA	
c) SSRI d) Combination of B and C	
,	_
Answ	er: A
57- Drug that does not use in Alzheimer disease	
TCA	
re. Not a side effect of an entiroxychotics.	
58- Not a side effect of an antipsychotics:	
hypotonia	
59- A patient has been hearing voices talking about him over the last three weeks and he can't see them, which	
psychopathology:	

Answer: Disorders of Perception. 60- A patient when asked about his level of education, talked about his school, his friends and concluded by saying, " that was the end of my schooling". Which demonstrates his answer: **Answer: Circumstantiality** 61- Increasing the amount of Heroin to achieve the same initial effect: **Answer: Tolerance** 62- Occurs in REM sleep: Answer: Loss of muscle tone 63- Which is antipsychotic: Answer: Quetiapine 64- A student who is afraid to give seminars as he is sensitive to negative evaluation, most likely diagnosis is: Answer: Social phobia 65- Definition of Delusions: Answer: False belief that is held despite evidence of contrary 66- A patient presented to ER with severe respiratory depression, on physical examination he is unresponsive with a tattoo over a large area of his body, also there are needle marks on his forearm. Which drug is the most likely involved? Answer: Heroin (Doctor Radwan concluded that a large tattoo is sign of personality disorder or a substance abuse, and as the patient presented with severe respiratory depression that is a sign of heroin abuse) 67- A patient admitted after an RTA, was conscious and became confused after a few hours as he started wandering in the ward thinking he is home. The diagnosis: **Answer: Delirium** 68- A sign of seriousness of suicidal attempt: Answer: Suicidal note (A suicidal note means a planned suicidal attempt) 69- Hallucination is a disorder of:



Answer: Amenorrhea
82- Most important in delirium management:
Answer: Searching for underlying cause
83- True depression:
Answer: Physical symptoms are common presentation
84- True about Doctor-Patient relationship:
Answer: Medical schools should nurture patient centered relationship
85- Recurrent short episodes of depression that was treated. Three weeks late the patient started having symptoms
of mania, diagnosis:
Answer: Bipolar disorder
86- True about adjustment disorder:
Answer: Onset of symptoms within 3 months after trauma exposure
87- True about schizophrenia:
Answer: Dopamine over activity is thought to be the cause
88-A feature of REM sleep:
a) decrease respiratory rate.
b) decrease cerebral blood flow.
c) dreams
d) parasympathetic activation
*Answer: c) dreams
89- testimonial capacity is:
a) ability to testify in court.
b) capacity to write a will.
Answer: B) capacity to write a will
90- indicate seriousness of suicide:
a) using firearms
b) writing a will c) angry for being saved.
d) being a female

Answer: A&B both are correct, but according to the past B was the answer

```
91- Typical antipsychotic:
 Answer haloperidol
 92- first drug to give to extrapyramidal side effects:
A)olanzapine
B)diazepam
 C) procyclidine
 D)citalopram
 E) lithium CO3
                                                                                         *Answer is c) procyclidine
 92- the disease caused by decrease serotonin:
     a) depression
     b) schizophrenia
                                                                                           Answer is A) depression.
 93- hallucination best description:
 Answer is perception without external stimulus.
 94- used in alcohol dependence:
     a) Olanzapine
     b) Diazepam
                                                                                            Answer is B) diazepam.
 95- general anxiety disorder is:
     a) anxious in public
     b) anxious about everyday events
                                                                        Answer is b) anxious about everyday event.
 96-social phobia is:
 Fear of being embarrassed (fear of being criticized)
 97- case about someone who fears public, crowded places:
 Agoraphobia
 98- bipolar prevalence:
 1%
 99- ADHD:
     a) decrease dopamine.
```

b) decrease after the age of 15.	
	Answer A) decrease dopamine.
100- personality disorder wrong:	
Begin at the age of 18.	
101- reversible cause of dementia:	
a) Alzheimer disease	
b) pick's disease c) pseudodementia	
c) pseudodementia	<b>3</b>
and all arms described as a land all a serious also	Answer is c) pseudodementia.
102- allowed units of alcohol per week:	
20	
103- most common presentation of depression:	
physical symptoms	
10.4 not acceptated with payorlange	
104- not associated with narcolepsy:  Answer increased REM SLEEP	
Allswer uicreased Reivi Sleep	
105- Gender identity:	
36 months	
106-side effect of SSRI's:	
nausea	
107- Drug that does not cause dependence:	
olanzapine	
108- which of the following is characteristic for unipolar depression:	
Anhedonia	
109- paranoid delusions are found in all of the following except:	
a)Depression	
b)Schizophrenia	
c)OCD	
d)Dementia	
e) Substance abuse	

	Answer) OCD * (Paranoid Delusions are not found in OCD)
110- Thought Echo is:	
a)A form of thought broadcasting	
b)Abnormal thought content	
c)An illusion	
d)Idea of reference	
,	
e) A hallucinatory experience	
	Answer: A
111- which of the following drugs causes dryness of mout	n the most:
a) Risperidone	
b) Fluoxetine	
c) Chloropyramine	
	Answer Chloropyramine * (antihistamine)
112- which of the following statements is true regarding l	Dr-Pt relationship:
A. It doesn't affect the course of the health problem	n.
B. Med schools should nurture patient centered me	edicine.
C. In the relationship the pt. has no active role to p	lay
D. Should concentrate on the disease not the patien	its' emotions.
	Answer: B
113- unlikely feature of neuroleptic malignant syndrome	
Leucopenia (true is leukocytosis)	
114- which of the following disturbances is associated with	h highest risk of suicide
a) Paranoid schizophrenia	
b) Dissociative disorder	
c) Major depression * (Bipolar is also high)	
d) Dementia	
	Answer: C Major depression * (Bipolar is also high)
115- all of the following are impulse control disorders exc	
·	~~~
conversion disorder (not impulse control disorder)	

Answer: E

117- which of the following is a feature of schizoid personality

a) Somatizationb) Dissociative

e) Malingering

c) Hypochondriasis

d) Munchhausen syndrome

116- unconscious mechanisms play a role in all of the following except:

a)	Desire to mix with others.
ь)	Anxiety
c)	Acceptance of abnormal traits
d)	Self-Importance
e)	Dependency
•	Answer: D
_	
_	came to the ER saying that for the last few months he has been hearing voices talking to him, when he was
asked i	he was sure, he said 100%, this shows which psychopathology:
a)	Derealization
ь)	Disorder of thought content
c)	Disorder of perception
	Answer: C
119- dia	gnosis of all of the following occurs in childhood or adolescents except
,	Separation anxiety
a) b)	Conduct disorder.
b)	ODD
c) d)	Personality disorder (only occurs after adulthood)
۵	resonancy disorder (only occurs areas additiood)
	Answer: D
120- CB	Γ effective in all except:
a)	Panic disorder
ь)	PTSD
c)	Social anxiety
d)	Depression
<b>e</b> )	Antisocial PD
	Answer: E
121- whi	ch of the following is associated with repeated overdose
a)	Munchhausen syndrome
b)	OCD
c)	Paranoid personality disorder
<b>d</b> )	Borderline PD
	Answer: D
-00	
122- WIII	ch of the following drugs associated with highest mortality among users:
Heroin	
123- whi	ch is true of depression:
a)	Physical symptoms are a common presentation.
ь)	Affects 2% of the general population.
c)	5% commit suicide
	Answer: A
12 <i>4-</i> Dei	personalization:
	<del></del>

Feeling	g of unreal self	
125- wh	nich of the following is a sign of seriousness in a suicidal attempt:	
suicida	al note	
126- wh	nich of the following is true regarding bipolar disorder:	
a)	Only presents in middle age.	
ь)	No genetic component	
c)	Treated by lithium.	
		Answer C
127- wh	tich of the following is a reversible dementia	
a)	Alzheimer's	
b)	Picks disease	
c)	CÌD	
d)	Alcohol induced.	
	,z	Answer D
128- which o	of the following alerts the doctor for the possibility of alcohol dependence when examining a patien	ıt:
a)	High blood pressure	
-) b)	Excessive cigarette smoking	
c)	Peripheral neuropathy	
,		Answer C
-		Miswer C
129- abi	normality of which system mostly associated with schizophrenia	
Dopaminergic		
130- wh	nich is a sleep disorder	
a)	Night Vision	
ь́)	Panic disorder	
c)	Cataplexy	
d)	Kleinn Levin disorder	
e)	Cautard syndrome	
		Answer D
131- whi	ich of the following is associated with anorexia nervosa	
a)	Amenorrhea	
ь)	Normal body image	
<b>c</b> )	Low blood pressure	
d)	Body weight 20% above normal	
	A	Answer A

132- mo	st imp in Delirium management:
a)	V fluids
<b>b</b> )	Search for the underlying cause
c)	Low light in the room
	Answer B
177-2-2	chizophrenic mother wants to get pregnant and asks about the possibility of having a schizophrenic child:
12% (ris	k of schizophrenic offspring for schizophrenic mother)
134- an	example of behavioral therapy:
a)	Gestalt therapy
ь)	Hypnosis
c)	Modeling
d)	Psychoanalysis
	Answer C
135- wh	ich is true regarding psychotherapy:
a)	Has no significant effect on brain neurotransmitters
-) b)	Aims to improve client functioning.
c)	Not suitable for children below 12 yrs.
,	Answer B
l	
	aracteristic for paranoid PD:
a)	Reckless behavior
ь)	Magical thinking
c)	Mistrust
	Answer C
13 <b>7-</b> cha	racteristic of REM sleep
loss of	muscle tone
158- pat	ient PTA wakes up later confused agritated and thinks he is home:
_	ient RTA, wakes up later confused, agitated and thinks he is home:
Deliriu	m.
139- pat	ient with depressive symptoms prescribed antidepressants, 3 weeks later presents with agitation, lack of
sleep, (	manic features), what did the doctor miss:
Bipolar	disorder
140- inc	crease suspicion to organic cause of acute psychosis:
,	
a)	High blood pressure

ь)	Peripheral neuropathy	
<b>c</b> )	Confusion	
d)	Auditory hallucination	
	Answer B	
141- a p	t. presents to the ER with Resp Depression, pinpoint pupils, and needle marks with Tattoo, most likely drug	
Heroin		
142- DV	romania is:	
an unp	oulse to set fires.	
143- ge	nder identity is established at what ages?	
<b>3</b> 6 moi	nths	
<b>J</b>		
144- A	characteristic of Schizoid personality disorder:	
a)	importance	
ь)	anxiety	
<b>c</b> )	accept abnormal traits.	
	Answer A	
145- Sw	oconscious mechanism one wrong:	
145-04	·	
a)	Malingering	
ь)	Dissociative	
c)	Somatization	
	Answer A	
146 <b>-</b> Pe	rsecutory Delusion not in:	
a)	Depression	
ь́)	OCD	
,	Answer: b	
147- inc	rease suspicion of organic cause to acute psychosis:	
a)	High blood pressure	
ь)	Confusion	
<b>c</b> )	Peripheral neuropathy	
<b>d</b> )	Auditory hallucination	
	Answer is c.	
148- sle	ep disorder:	
	n due syndrome.	
B-Cata	olexy	
	Answer B	
149 <b>-</b> Th	ought echo:	

B-Audi	tory hallucination	
		Answer B
150- A c	characteristic of unipolar depression	
Low sel	lf-esteem.	
iei- One	e is a behavioral therapy technique	
Floodin		
152- Per	rsecutory delusions in all except	
OCD		
153- Dr-	Patient relationships; what is true	
	alistic relationship with the patient is unacceptable.	
	nich is not impulse disorder	
Conver	sion disorder.	
155- All	disorders have unconscious mechanisms except	
a)	Münchhausen (we chose this,	
<b>b</b> )	Somatization	
<b>c</b> )	Hypochondriasis	
d)	Dissociative Symptoms	
	Answer: A was chosen malingering was not a choice	
156- Wh	ich disorder is associated with repeated drug overdose	
a)	Borderline	
ь)	Münchhausen	
<b>c</b> )	Depression	
		ANSWER A
157- Coş	gnitive behavioral therapy effective in all except	
a)	Panic disorder	
ь́)	PTSD	
c)	Depression	
<b>d</b> )	Anxiety	
<b>e</b> )	Paranoid schizophrenia	
		A NICWED E

A-Broadcasting thought

158- Which of the following occurs in REM  $\,$ 

a) Night terrors

Night terrors and enuresis are not typically associated with	REM sleep.
159- True about Schizoid personality	
Solitary behavior.	
160- True about gender identity disorder	
A-Difficult to treat.	
B-More common in west civilization	
	ANSWER: A
161- Risk for bipolar mother having bipolar child	
20% (according to Medscape 15-25%)	
options were.	
10,	
30, and higher	
162- Sign of alcohol dependence	
A-Liver enlargement	
B-Vitamin B6 deficiency	
	ANSWER: A
163- Associated with anorexia	
A-Anxiety	
B-Weakness	
	ANSWER: A
164- Increased suicide attempt seriousness in?	
<ul><li>A. Being female (no, being male has a higher risk)</li><li>B. Being married (no, it DECREASES the risk of serious attempt)</li></ul>	
C. Previous attempts (correct answer as stated by Dr Laith Abbadi)	
D. Being depressed	
165- Sign of organic underlying cause for psychosis	
Upper limb paresis.	
ACCORDER 1 PR A 1	
166- CBT is ineffective in:	
Paranoid Schizophrenia	
(as checked by Dr Laith Abbadi)	

b) Enuresis

Monoamine System			
168- Associated with anorexia nervosa:			
A. Anxiety B. Weakness			
C. Normal body image			
D. Low blood pressure			
E. Body weight 20% above normal			
A	Inswer: a		
169- cause agranulocytosis:			
clozapine			
170- Dysmorphophobia, one is true:			
seeks surgery			
tet velikaly to happen in esting disarder.			
171- unlikely to happen in eating disorder:			
hyperkalemia			
172- considered hallmark in Alzheimer:			
amyloid plaques			
173- not an indicator for true hallucination:			
Voices inside the head			
174- unlikely in paranoid type of schizophrenia:			
waxy flexibility			
175- occur in conversion disorder:			
unconscious anxiety			
unconscious anxiety			
176- true about agoraphobia:			
comorbid major depression			
177- true about bipolar:			
peak of onset in late adolescent and early adulthood			

167- Abnormalities in which system is best associated with depression?

178- true about oppositional defiant disorder:	
seeking revenge	
179- true about Asperger:	
normal cognitive development	
180- shared side effect between TCA and SSRI:	
sexual dysfunction	
181- unlikely to cause hyperprolactinemia:	
alprazolam	
182- good predictor for schizophrenia:	
previous history of mood disorder	
previous history of moor disorder	
10- whysical factors of DD7 dependence	
183- physical feature of BDZ dependence:	
tremor	
of a complete block on the delication	
184- carries highest suicidal risk:	
a) bipolar b) depression	
·,	Answer is A
185- true about panic disorder:	
anticipatory anxiety	
unctofputor, unknow,	
186- true about tardive dyskinesia:	
incurable	
187- true about homosexuality:	
feature of sexual orientation	
ANADAR O DE SOMME OF FORESTORE	
188- untrue about cognitive behavioral thereny	
188- untrue about cognitive behavioral therapy:	
exploration of unconsciousness	
-0	
189-cognitive test used in dementia:	
mini mental state examination	

```
190- not a cocaine withdrawal symptom:
elation
191- false about depression in the elderly:
lower suicidal rate
192- not a risk factor for suicide:
childless marriage
193-. one of these is a paraphylia:
fitishism.
194- one of these drugs is lethal in overdose:
amitriptyline
195- a male seeing snakes crawling on the wall, one of these is true:
an optimal dose of risperidone is appropriate treatment.
196- false regarding schizophrenia:
females have an earlier onset of the dis than males
197- one of these is an indication of clonazipine:
persistent negative symptoms after being treated for 6 weeks with 3 antipsychotics at optimal dose.
199- least likely diagnosis in a 16 yr. old child with ADHD:
antisocial personality disorder
200-. to dx bipolar disorder you need symptoms for:
2 weeks.
201- factitious disordered:
unconscious of the motivation
202- true about autism:
```

poorly developed non-verbal social communication

203- false regarding anorexia nervosa and bulimia nervosa:
bulimia has an earlier onset
204- treatment of specific phobia:
flooding
205- contraindication for ECT:
brain tumor
206- true regarding paranoid delusional disorder:
preserved social life and function
207-Least fatal in OD:
Correct answer: alprazolam
108 Chance wrong angewer
208- Choose wrong answer:
anticonvulsant drugs used routinely in management of alcohol withdrawal
209- Cognitive behavioral therapy is least effective for:
Answer: antisocial personality disorder (ASP) I chose agoraphobia.
210- NT least incriminated in depression:
a) GABA
b) NE
c) Ach d) DA
Answer is C
211- Wrong answer:
less than 1/3 of children are dry at age 5 (Qabout enuresis) Typically, most children are bowel and bladder trained
by age 3 years.
212- True about OCD:
Compulsions must be preceded by intrusive thoughts obsessions can be without compulsions, but compulsions
can't be without obsessions.
213- Alzheimer:

```
higher prevalence in > 80 y old
214- trichotillomania:
impulse control disses
215- wrong:
eating dis/anorexia more prevalent in lower economic class
216- Autonomic arousal/sympathetic overactivity present in all except:
    a) conversion disorder
    b) GAD
    c) PTSD
                                                                                                        Answer is A
217- exposure and response prevention Rx of:
OCD
218-3 episodes of depression: Tx should continue for how long?
answer: for life
(Dr explained: 1 depressive episode: Rx for 1 y; 2nd episode: for 2 y; 3 or more episodes: for life)
219- Most effective Rx of panic dis:
fluoxetine
220- Bad dreams least likely with:
mania
221- Thought disorder in depression:
negative thoughts
222- wrong about manic depression:
minimal genetic role
223- Ass with metabolic syndrome:
clozapine (other options were not antipsychotics)
224- true about conduct disorder:
```

	predisposes to	antisocial personality disorder in adulthood	
	225- True about	PTSD:	
	startle response	е	
	226-True about	GAD:	
	stress for no su	fficient reason	
	22 <b>7-</b> Which of th	he following most likely to cause psychosis?	
	a)	Cocaine	
	b)	Cannabis	
			Answer is a
	228- Which of th	he following is an effect of the drugs causing addiction?	
	a)	increase serotonin.	
	ь)	decrease noradrenaline.	
	<b>c</b> )	causing tolerance	
	d)	decrease dopamine.	
			Answer is c
	229- Which of the	hese is true about panic disorder?	
a)	diagnosed fron	n single episode.	
<b>b</b> )	occurs in femal	es more than males	
<b>c</b> )	they will not be	enefit from cognitive therapy.	
			Answer is b
	230- Which of the	hese is not true about personality disorders?	
Pat	cients with perso	nality disorders usually have low IQ ***	
		ne following is a feature of NON-REM?	
It c	comprises 75% of	the sleep cycle ***	
	232- Which of the	he following alcohol blood levels will cause abnormal behavior?	
	a) 20 mg		
	b) 40 mg		
	c) 50 mg		
	d) 100 mg		
	e) 200 mg	7	
			Answer is E
	233- Which of the	he following increase the risk of suicide?	
	a) being f	female	

b) being physician

Answer is C

	234	- Which of the following drugs is anti-depressant?	
	a)	Alprazolam	
	<b>b</b> )	citalopram	
			Answer is B
	0	Which of the following in not a part of mental examination?	
	<b>4</b> 35	•	
	a)	speech	
	ь)	mood	
	c)	neurological examination	
	d)	memory	
	<b>e</b> )	judgment	Answer is c
			Allswer is C
	236-	Which of the following defines hallucinations?	
	a)	perception without an external stimulus	
	<b>b</b> )	An image presented to consciousness.	
	<b>c</b> )	perceived things in a different way than normal	
	d)	misinterpretation of external stimulus	
			Answer is a
	237-	Which of the following statements is true about schizophrenia?	
	a)	mostly occurs in the middle age.	
	<b>ь</b> )	genetic is the most common etiology	
			Answer is b
	- 0	N. C.J. C.D	
	238-	· All of the following indicate seriousness of suicide except?	
	<b>a</b> )	previous attempt	
	ь)	being married	
			Answer is b
	239-	· Which of the following drugs may cause addiction?	
λln		lam ***	
7 X P	1 020		
	240	- The definition of Stupor?	
Cor	ıscio	us but immobile ***	
	241-	Hypomania is a disorder of?	
	•	a) cognition	
		b) memory	
		c) mood	

242- Risk of suicide in depression?

a)	5%	
b)	10%	
c)	15%	
<b>d</b> )	20%	
e)	25%	
		Answer is b
•	Wilter 6 d. 6 H	
24	3- Which of the following is a common lithium side effect?	
<b>a</b> )	Nausea	
ь)	polyurea	
		Answer is b
24.	4- Which of the following indicates that there is an organic disease?	
altered	l level of consciousness ***	
24	5- Underactivity of Ach will cause?	
	a) schizophrenia	
	b) anxiety	
	c) dementia	
		Answer is c
24	6- Which of the following statements is true about depression?	
a)	It is a leading cause of disability in community	
	•	
b)	35% of general populations have depression.	
b) с)	35% of general populations have depression.  females are four time to be affected than males.	
c)	35% of general populations have depression.  females are four time to be affected than males.	Anewerica
c)	females are four time to be affected than males.	Answer is a
c)		Answer is a
c) 24	females are four time to be affected than males.	Answer is a
c) 24	females are four time to be affected than males. 7- Which of the following drugs is anti-psychotic?	Answer is a
c) 24 risperi	females are four time to be affected than males. 7- Which of the following drugs is anti-psychotic?	Answer is a
c) 24 risperi	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***	Answer is a
c) 24 risperi	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?	Answer is a
c) 24 risperi	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations	Answer is a
c) 24 risperi	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal	Answer is a
c) 24 risperi	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion	Answer is a
c) 24 risperi	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion  d) passivity phenomena	
24 risperi 24	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion  d) passivity phenomena	
24/risperi 24/ 24/ a)	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion  d) passivity phenomena  9- Which of the following drugs will treat extrapyramidal symptoms?  benzhexol	
24 risperi 24	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion  d) passivity phenomena	
24/risperi 24/ 24/ a)	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion  d) passivity phenomena  9- Which of the following drugs will treat extrapyramidal symptoms?  benzhexol	
24/risperi 24/a)	females are four time to be affected than males.  7- Which of the following drugs is anti-psychotic?  done ***  8- Which of the following is a negative symptom?  a) hallucinations  b) social withdrawal  c) delusion  d) passivity phenomena  9- Which of the following drugs will treat extrapyramidal symptoms?  benzhexol	Answer is b

b) Anxiety disorders are risk factors of low achievement Answer is b 251- Case (male experienced a panic attack then he became afraid of getting out from his home alone) Agoraphobia \*\*\* 252- Generalized anxiety disorder? Anxiety about everyday issues \*\*\* 253- Social phobia is which of the following? phobia from crowded places phobia from negative evaluation Answer is B 254- overactivity of dopamine will cause? schizophrenia depression Answer is a 255- Which of the following is a good prognostic factor of schizophrenia? a) insidious onset b) family history c) precipitating factors \*\*\* d) negative symptoms Answer is c 256- Which of the following will cause irreversible dementia? brain tumor dementia arteriosclerotic dementia \*\*\* hypothyroidism dementia Answer is b 257- Which of the following occurs in dementia? Fluctuation in consciousness is a primary mechanicm Hallucinations and delusions are prominent symptoms The onset is usually acute D. All cases are untreatable Amnesia for names are experienced early in the course of the disease ANSWER: A 258- In people over the age of 65, one of the following is True: A. Parasuicide is more common than suicide About 20 percent have dementia

- Severe memory loss is common D. Anxiety disorders are more common than depressive disorders E. Any grade of amnesia indicates Alzheimer disease ANSWER B 259- Which of the following is True of adjustment disorder? A. Is a response to a serious unexpected traumatic event The symptoms usually last more than six months
- - Psychosocial performance is unaffected
  - D. Conduct symptoms are the frequent presentation in adolescents with this disorder
  - Grief is an ..... disorder

ANSWER A

260-Which of the following is True of Impulse Control Disorder?

- A. Fugue can be a presenting feature.
- B. May present with trichotillomania.
- C. May present with unexplained physical symptoms.
- D. Includes malingering.
- Is associated with experiencing depersonalization.

Answer is B

261-Which of the following is a characteristic feature of Post-Traumatic Stress Disorder?

- Involvement in conversations related to the trauma.
- Recurrent distressing dreams
- Frequently visiting the place where the trauma occurred.
- Lowered arousal level
- The patient is always a survived victim of the trauma.

Answer is B

- 262. Desensitization is effective in the treatment of:
- A Specific phobia
- B. Social phobia
- C. Agoraphobia
- D. Obsessive compulsive disorder
- E. Conversion disorder

ANSWER is A

263. Which of the following medications would be the most appropriate to prescribe for a patient with Bipolar Depression?

- A. Haloperidol
- **Bromazepam**
- Valproic acid C.
- Fluoxetine D.
- Risperidone

Answer is C

264. Acute drug induced dystonia can be due to the use of:	
A. Procyclidine	
B. Alprazolam	
C. Propranolol	
D. Olanzapine	
E. Haloperidol	
	Answer is E
265. Which of the following is a REM sleep disorder?	
A. Nightmare	
B. Sleep walking	
C. Sleep talking	
D. Jet lag	
E. Enuresis	
	Answer is A
266. Dryness of the mouth is a recognized adverse effect of one of	
A. Fluoxetine	
B. Imipramine	
C. Diazepam	
D. Velafaxine	
E. Olanzapine	
	Answer is E
267. Which of the following is true of treatment with lithium?	
A. Lithium is the treatment of choice in schizophrenia.	
B. On long term use causes hyperthyroidism  C. Hos acceptated with marked antichelineratic adverse effects	
C. Use associated with marked anticholinergic adverse effects.	
D. Useful in the prophylactic treatment of bipolar  E. Safa to you in programs patient	
E. Safe to use in pregnant patient.	Answer is D
269. Which of the following statements concerning behavior thereasy is true?	Allswer is D
268. Which of the following statements concerning behavior therapy is true?	
A. Behavior therapy is more effective than SSRIs in the treatment of depressive disorder.	
B. Includes thought stopping technique in OCD.	
C. Less effective than benzodiazepines in the treatment of agoraphobia	
D. Based on the exploration of the unconscious.	
E. Is a form of talking therapy.	
	Answer is E
269. Which of the following is true of autistic disorder?	
A. Basically, it is a disorder of intelligence.	
B. Distractibility and short attention span are pathognomonic diagnostic features.	
C. Onset is often before the age of 3.	
D. Is a form of schizophrenia of childhood.	
E. Response to antipsychotic medications	
	Answer is C
270. Which of the following is true of bipolar disorder (manic type)?	
A. Affects young girls more than men.	

B. At least two weeks of symptoms required for the diagnosis.	
C. Manic episode always immediately followed by a depressive episode.	
D. Usually associated with third-person auditory hallucinations.	
E. Mood stabilizers are preferred over antipsychotics in the long-term treatment.	
	Answer is E
271. Which of the following drugs is most useful in the treatment of ADHD?	
A. Haloperidol	
B. Fluoxetine	
C. Diazepam	
D. Methyl phenidate	
E. Riperadon	
	Answer is D
272. Which of the following symptoms is the most diagnostic of schizophrenia?	
A-Affective flattening	
B-Persecutory ideas	
C-Overvalued ideas	
D Primary delusion	
E. Grandiose delusion	
	Answer is D
273. Which of the following medications is most useful in the treatment of a patient with major	r depressive episode?
A-Chlorpromazine	
B Fluoxetine.	
C. Carbamazepine	
D-Olanzapine	
E-Risperidone	
	Answer is B
On a Which of the following is true of shildhood system?	
274. Which of the following is true of childhood autism?	
A. Is associated with conduct disorder in childhood.	
B. Autistic features appear after the age of 3 years.	
C. Equally prevalent in male and female children	
D. Can be effectively treated by atypical antipsychotics.  E. Social and communication skills are abnormally developed.	
E. Social and communication skills are abnormally developed.	Answer is E
275. Dissociative disorder includes one of the following:	Answer to L
A. Acute paraplegia	
B. Fugue	
C. Dysmorphophobia	
D. Hypochondriasis	
E. Dysthymia	
	Answer is B
276. Which of the following is an effective medication in the management of panic disorder?	

A.	Risperidone	
B.	Haloperidol	
C.	Fluoxetine	
D.	Sodium valproate	
E.	Clozapine	
	•	Answer is C
277.	Which of the following is a diagnostic feature of Somatization disorder?	
_	·	
A. -	Weight loss of 25% below expected body mass.	
В.	Increased libido	
C.	Preoccupation with physical appearance	
D.	Multiple physical complaints without organic pathology for more than 2 years	
E.	Repeated investigations with normal results reassure the patient.	
		Answer is D
2 <del>7</del> 8.	In the management of major depressive episode one of the following statements is true:	
A.	In severe depression SSRI's give more rapid therapeutic effect than ECT.	
B.	Lithium is superior to TCA agents in treatment of bipolar depression.	
C.	Prolonged regular use of benzodiazepines is considered safe.	
D.	In the presence of ischemic heart disease, it is safe to prescribe TCA agents.	
E.	ECT is contraindicated if the patient is an elderly.	
		Answer is B
279.	Which of the following is true of ADHD?	
A.	Characterized by hyperactivity in one setting.	
В.	Distractibility and short attention span are prominent features.	
Б. С.		
_	Methylphenidate can be instituted at age of 4 years.  Never seen in adults.	
D.		
E.	Benzodiazepine drugs are effective.	America D
<b>.</b> 0	With a field of the field of the control of the con	Answer is B
280	Which of the following medications may cause hypothyroidism?	
A.	Lithium carbonate	
В.	Haloperidol	
C.	Valproate	
D.	Benzhexol	
E.	Methylphenidate	
		Answer is A
281.	Hallucinations are unlikely to be experienced in one of the following conditions:	
A.	Schizoaffective disorder	
В.	Bipolar disorder	
C.	Major depressive episode	
D.	Delusional disorder	
E.	Substance use disorder.	
		Answer is d
282-	A patient presented to the clinic with acute drug induced dystonia, which of the following	
	ely to help?	middle and a middle
A.	Procyclidine	

B. Cholordiceoxide	
C. Propranolol	
D. Olanzapine-	
E. Sertraline	
	Answer is a
00_ With a fair fall and a second of the sec	
283- Which of the following symptoms indicate alcohol dependency?	
A. Morning shakes	
B. Morning hangover	
C. Maming dyspepsia	
D. Morning headache	
E. Morning blackouts	
	Answer is B
284. Which of the following conditions is associated with short REM sleep latency?	
A Generalized anxiety disorder.	
B. Obsessive compulsive disorder	
C.Narcolepsy depression sleep dep	
D. Schizophrenia	
E. Mania	
	Answer is C
285. Which of the following is a benzodiazepine?	
A. Citalopram	
B. Alprazolam	
C.Escitalopram	
D.Gabapentin	
E. Disulfiram	
	Answer is B
286.All the following regarding cannabis are true, EXCEPT:	
A. Tetrahydrocannabinol is the active substance.	
B. Cannabis smoke contains the same carcinogens as tobacco.	
<ul><li>C. May be detectable in the urine up to 4 weeks after use.</li><li>D. It is a liquid-soluble substance.</li></ul>	
E. It belongs to the opioids class of drugs.	
	Answer is E
28x Which of the following is true about premenetrical describerio disorder?	
287. Which of the following is true about premenstrual dysphoric disorder?	
A. Symptoms occur between ovulation and menstruation.	
B. Symptoms occur at any phase of menstrual cycle.	
C. Considered normal when occurs in girls below 18 years of age.	
D. Has a weak association with depression  E. Occurs in less than 20% of women.	
2. Cooled the took likely 20/0 of wolliely	Answer is A

288.Which of the following is unlikely to be associated with depression?	
A. Hypothyroidism	
B. Addison's disease	
C. Pheochromocytoma.	
D. Pellagra	
E. Cushing's disease:	
	Answer is C
289. Which of the following drugs may cause depression?	
A. Lithium carbonate	
B. Imipramine.	
C. Haloperidol	
D. Fluoxetine.	
E. Valproate	
	Answer is C
290. Which of the following is more diagnostic of schizophrenia?	
A Persecutory delusions	
B. Obsessional ruminations	
C.Command hallucinations	
D. Blunt effect	
E. Pressure of speech	
	Answer is A
291. Which of the following is a good prognostic sign in schizophrenia?	
A. Family history of schizophrenia  B. Acute onset of illness	
C. Poor affective response Å	
D. High expressed emotion family environment	
E. Onset before the age of 18 years	
	Answer is B
292. All the following are correct regarding schizophrenia, EXCEPT:	
A. Both schizophrenia and affective symptoms are present at the same time.	
B. The prognosis is usually better than that of bipolar.	
C. Younger people tend to have associated mania while older people have depressive. symptoms	
D. History of schizophrenia occurs in excess of schizophrenia in their relatives.	
•	
E. Treatment consists of antipsychotics and mood stabilizers.	
	Answer is B
293. Which of the following is true regarding puerperal psychosis?	
A. Relapses are usually absent.	
B. Usually resistant to treatment	
C.Usually of acute onset	
D. Response to ECT is poor.	
E. More common with last born baby.	
EVALUE COMMINIST WEEK LEED COMMINISTER	

	Answer is C
294. All the following are true regarding Down syndrome, EXCEPT:	
A. Is the commonest cause of severe learning disability.	
B. The majority of affected patients have trisomy 21.	
C. Patients suffer a wide range of physical abnormalities.	
D Patients have normal fertility.	
E. Risk to mothers having a baby with Down syndrome increases with age.	
	Answer is D
	ARISWOI IS D
295. All the following are correct regarding anorexia nervosa, EXCEPT:	
A. Low level of thyroid hormones	
B. Usually develops leukocytosis.	
C.Disturbance of the way in which the body is perceived.	
D. Occurs mainly in young women.	
E. Depression is common.	
·	Answer is B
	Aniswer to B
296. All the following are somatic symptoms of anxiety, EXCEPT:	
A. Palpitations	
B. Hyperventilation	
C.Depersonalization	
D. Urinary frequency	
E. Nausea	
	Answer is D
	THIS WOT US D
297. All the following point to good practice in using benzodiazepines, EXCEPT:	
A. Short term use	
B. Lowest effective dose	
C.Use large dose in primary insomnia	
D. Use to deal with specific problems.	
E. Intermittent doses	
	Answer is C
	AMISWEL IS C
298. Which of the following disorders is characterized by impaired reality testing?	
A. OCD	
B. Delusional disorder	
C. Dissociative disorder	
D. Somatization disorder	
E. Impulsive control disorder	
	Answer is B
299. All the following occur in bulimia nervosa, EXCEPT:	
A. Parotid gland enlargement	

B. Disordered body image

C. Family conflict

D. Dental erosion E. Body weight less than 90% of normal Answer is E 300. Features of depression include any of the following, EXCEPT: A. Fatigue B. Lack of interest C. Psychomotor retardation D.Mood incongruent hallucinations E. Anhedonia Answer is D 301. All of the following are true of agoraphobia, EXCEPT Occurs more in females than males. Affects children more than adults. May be associated with panic symptoms. Better treated by flooding than desensitization. Behavioral therapy is better than drug therapy. Answer is B 302. Supportive psychotherapy involves all the following, EXCEPT: A. Listening to the patient Giving guidance to the patient Analytic. C.Exploration of the unconscious conflicts D. Can be practiced by doctors in all specialties. E. Useful for patients with mental and physical disorders Answer is C 303. All of the following are true of schizophrenia, EXCEPT: Long term administration of antipsychotic medications is mandatory. Living in an environment with high expressed emotions results in poor outcome. Genetic factors play a role in causation. Negative symptoms respond equally well to typical and atypical antipsychotics. Presence of schizoid premorbid personality predicts poor prognosis. Answer is D 304. OCD is characterized by: A. Thoughts that are inserted by external agencies. Thoughts that are usually can be easily resisted. C.Thoughts that are intrusive and considered by the patient as alien. D.Thoughts that are meaningful. E. Thoughts that induce pleasure Answer is C 305. Labile affect is: A. A disorder of formal thought disorder

Means the patient laughs a minute and cries the next for no obvious reason. В. C. A bipolar disorder syndrome A symptom of dysthymia D. A term which is synonymous with anhedonia Answer is B 306. Which of the following is a convenient therapeutic approach for Individuals with antisocial personality disorder? A. Cognitive behavioral therapy Therapeutic community C. **ECT** Psychodynamic psychotherapy D. Antipsychotic medication Answer is B 307. Which of the following is a positive schizophrenic feature? Ambivalence **Blunted affect** Catatonic symptoms D.Command hallucination E. Ideas of reference Answer is D 308. Which of the following occurs in bulimia nervosa? A. Body mass index reduction of more than 15% of normal Distorted body image В. C. Lack of appetite Dieting D. Amenorrhea Answer is B 309. Which of the following medications is effective in the treatment of schizophrenia? Imipramine B. Fluoxetine C.Risperidone D. Lithium carbonate E. Valproate Answer is C 310. Which of the following is true of factitious disorder? A. Conversion symptoms are common. The patient is unconscious of the reason for producing the symptom. C. The patient is considered malingerer. D. The patient usually undergoes multiple surgical procedures. E. Munchasen syndrome is an example of this disorder. Answer is E

311. Which of the following differentiates delirium from other psychotic disorder	
A. Ideas of reference	
B. Overvalued ideas	
C. Voices of neighbors heard both by the patient and others in his voice.	
D. Visual hallucinations	
E. Clouded consciousness	
	Answer is E
312. All the following are recognized therapeutic approaches in psychiatry, EXCEPT:	
A. Cognitive behavioral therapy	
B. Interpersonal therapy	
C. Academic education	
D. Faith healing	
E. Pharmacotherapy	
•	Answer is D
	Miswer ts D
313. Regarding the etiology of mental disorders, one of the following is true:	
A. Migration	
B Dietary regulation	
C. Marriage	
D. Wealth	
E. Long employment	
	Answer is A
314. Completed suicide is more likely to occur in:	
A. middle-aged men	
B. young age women	
C. adolescents of both sexes	
D. Patients with dysthymia	
E. Hysterical personality disorder	
	Answer is A
315. Which of the following is an impulse control disorder?	
A. Erotomania	
B. Dysmorphophobia	
C. Trichotillomania	
D. Depersonalization	
E. Munchausen syndrome	
	Answer is C
316. One of the following is true of ADHD:	
A. onset before nine years old	
B. fifty percent co morbidity with autism	
C. Male to female ratio is 1 to 3.	
D. Short attention span is essential for the diagnosis.	
E. Only 5% have conduct disorder feature.	

	Answer is A
317. All the following features are true childhood autism, EXCEPT:	
A. Qualitative impairment in social interaction	
B. Insistence in repeating the same.	
C. significant delay in language development	
D. Up to 5% develop epilepsy.	
E. Intelligence can be normal.	
	Answer is D
	TARISWOT US D
318.Which of the following is a conversion symptom?	
A. Depressed mood	
B. Fugue	
C. Aphonia	
D. Panic attack	
E. Depersonalization	
319. Which of the following is an example of third-person hallucination?	Answer is C
A. A voice inside the head saying to the patient "you are useless."	
B. A voice giving orders to the patient not heard by others close to him.	
C. Voices of neighbors heard both by the patient and others in his vicinity.	
D. A voice commenting on the patient's actions.	
E. Voices of people conversating and referring to the patient	
	Answer is E
320. Which of the following is a feature of cocaine dependence?	
A. Sedation	
B. Hypervigilance	
C. Reduced perceptual sensitivity.	
D. Pinpoint pupils	
E. Hypoacusis	
	Answer is B
321. Which of the following is the drug of choice in treatment of unipolar depression?	
A. Haloperidol	
B. Bromazepam	
C. Beta blockers	

Answer is D

D.Citalopram

B.

C.

E. Lithium carbonate

322. Which of the following with regard to alcohol withdrawal features is true?

A. Delirium tremens is expected to occur 72 hours after the last drink. Alcohol is advised as a best action to abort withdrawal symptoms.

They respond dramatically to anticonvulsant medications.

Auditory hallucinations are predominant withdrawal feature.

Memory blackouts indicate alcohol dependence.

323. Which of the following is true of bipolar disorder?	
A. Thought echo is common.	
B. Overvalued ideas are characteristic.	
C. Anhedonia is frequently complained of	
D.Pressure of speech is remarkable.	
E. Command hallucination are frequent.	
	Answer is D
324. which of the following psychotropic medications is likely to be fatal in overdose?	
A. imipramine	
B. Fluoxetine	
C Alprazolam	
D-Fluvoxamine	
E-Olanzapine	
	Answer is A
325. Panic attacks occur in all the following conditions, EXCEPT:	
A-Substance withdrawal	
B. Lithium toxicity	
C- Phaeochromocytoma	
D-Hyperparathyroidism	
E-stress disorders	
	Answer is E
326. Which of the following is a dynamic psychotherapy term?	
A. Desensitization	
B. Reinforcement	
C. Free association	
D. Exposure	
E. Response prevention	
	Answer is C
327. One of the following does not differentiate grief from depression:	
A. Severe suicidal ideas	
B. prominent delusion of guilt	
C. Response to antidepressant therapy	
D. sadness	
E. Anhedonia	
	Answer is D
328. Erectile dysfunction is caused by which one of the following?	

A. SSRIs

Answer is A

B. Sildenafil	
C-Mania	
D-Testosterone	
E-Yohimbine	
	Answer is A
	Miswer is A
329-Which of the following statements is true about anorexia nervosa?	
A. Behavior that targets weight control	
B. Dieting that resulted in 20% body weight loss	
C. Depressed mood resulting from loss of weight.	
D. Use of appetizers to fight anorexia.	
E. Frequent psychiatric consultations	
	Answer is A
330. Which of the following statements is true about bulimia nervosa?	
A-The peak incidence occurs in boys between 15 and 19 years of age.	
B Dental caries is a consequence	
C-Constipation is a common feature.	
D-Increased prevalence in higher socioeconomic class	
E-Body mass index is below 17.5.	
	Answer is D
331. Which of the following statements is true regarding anxiety disorders?	
A-The fear is controllable.	
B-The fear can be reasoned away.	
C- Affects men more than women.	
D-Autonomic arousal is usually present.	
E-Libido is increased.	
	Answer is D
were Which of the following statements is Two of promature circulation?	
332. Which of the following statements is True of premature ejaculation?	
A. Response to behavior therapy is poor.	
B. May result from use of fluoxetine.	
C.May respond to squeeze technique.	
D. Is a form of Paraphilias.	
E. Is an adverse effect of SSRIs.	
	Answer is E
333. Which of the following statements is True of anorexia nervosa?	
A. The peak incidence occurs between 25 and 35 years of age.	
B. The female to male ratio is around 5:1.	
C. Diarrhea is a common feature.	
D. Affects only upper socioeconomic class.	
E. Body mass index is below 85%.	

Answer is B  334. Which of the following is True of Childhood Autism?  A. Social interaction improves by age 6.  B. Delayed language development is a marked feature.  C. LQ usually above 70.  D. Seizures affect about 75% of affected children.  E. Predominantly, is a disorder of male children.  Answer is B  335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25.  C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338. which of the following factors increases the risk of suicide in depression:  A. Penuale gender  B. Married  C. Old age  D. Yunner girls		
A. Social interaction improves by age 6.  B. Delayed language development is a marked feature.  C. I.Q. usually above 70.  D. Seizures affect about 75% of affected children.  E. Predominantly, is a disorder of male children.  Answer is B. 335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25.  C. Medication is the most effective treatment.  D. The major part in its ectology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C. 356. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B. 357. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C. 358. which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age		Answer is B
B. Delayed language development is a marked feature.  C. I.Q. usually above 70.  D. Seizures affect about 75% of affected children.  E. Predominantly, is a disorder of male children.  Answer is B  335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25.  C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338. which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	334. Which of the following is True of Childhood Autism?	
C. LQ usually above 70.  D. Seizures affect about 75% of affected children.  E. Predominantly, is a disorder of male children.  Answer is B  335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25.  C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	A. Social interaction improves by age 6.	
D. Seizures affect about 75% of affected children.  E. Predominantly, is a disorder of male children.  Answer is B  335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25. C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	B. Delayed language development is a marked feature.	
E. Predominantly, is a disorder of male children.  Answer is B  335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25.  C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	C. I.Q. usually above 70.	
335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25. C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	D. Seizures affect about 75% of affected children.	
335. Which of the following is True of Attention Deficit Hyperactivity Disorder?  A. Its prevalence in school-age children is in excess of 30%  B. All sufferers are free of the Disorder at age 25. C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	E. Predominantly, is a disorder of male children.	
A. Its prevalence in school-age children is in excess of 30% B. All sufferers are free of the Disorder at age 25. C. Medication is the most effective treatment. D. The major part in its etiology is a dominant gene. E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy? A. Very effective in treating a patient with schizophrenia. B. Uses thought diaries to identify thinking distortions. C. Effective treatment in specific phobia D. Usually, a long-term treatment E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia? A. Persecutory delusions B. Second person auditory hallucinations C. Thought echo. D. Illusions E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression: A. Female gender B. Married C. Old age		Answer is B
A. Its prevalence in school-age children is in excess of 30% B. All sufferers are free of the Disorder at age 25. C. Medication is the most effective treatment. D. The major part in its etiology is a dominant gene. E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy? A. Very effective in treating a patient with schizophrenia. B. Uses thought diaries to identify thinking distortions. C. Effective treatment in specific phobia D. Usually, a long-term treatment E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia? A. Persecutory delusions B. Second person auditory hallucinations C. Thought echo. D. Illusions E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression: A. Female gender B. Married C. Old age	775. Which of the following is True of Attention Deficit Hyperactivity Disorder?	
B. All sufferers are free of the Disorder at age 25. C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	•	
C. Medication is the most effective treatment.  D. The major part in its etiology is a dominant gene.  E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	- ·	
E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338. which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	•	
E. All of the sufferers have specific learning disabilities (dyslexia)  Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338. which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	D. The major part in its etiology is a dominant gene.	
Answer is C  336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age		
336. Which of the following is true of Cognitive Behavior Therapy?  A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	2. All of the surferers have spectic learning disabilities (dystexia)	
A. Very effective in treating a patient with schizophrenia.  B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338. which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age		Answer is C
B. Uses thought diaries to identify thinking distortions.  C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	336. Which of the following is true of Cognitive Behavior Therapy?	
C. Effective treatment in specific phobia  D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	A. Very effective in treating a patient with schizophrenia.	
D. Usually, a long-term treatment  E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	·	
E. Is a form of dynamic psychotherapy.  Answer is B  337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions  B. Second person auditory hallucinations  C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age		
Answer is B 337. Which of the followings is a characteristic positive symptom of schizophrenia?  A. Persecutory delusions B. Second person auditory hallucinations C. Thought echo. D. Illusions E. Obsessional symptoms  Answer is C 338.which of the following factors increases the risk of suicide in depression:  A. Female gender B. Married C. Old age	•	
A. Persecutory delusions B. Second person auditory hallucinations C. Thought echo. D. Illusions E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression: A. Female gender B. Married C. Old age	2. 10 1.0.11 of 2/1.111 po / 0.10 1.10 apr / 0.10 1.10 1.10 1.10 1.10 1.10 1.10 1.	Answer is B
<ul> <li>B. Second person auditory hallucinations</li> <li>C. Thought echo.</li> <li>D. Illusions</li> <li>E. Obsessional symptoms</li> <li>Answer is C</li> <li>338.which of the following factors increases the risk of suicide in depression:</li> <li>A. Female gender</li> <li>B. Married</li> <li>C. Old age</li> </ul>	337. Which of the followings is a characteristic positive symptom of schizophrenia?	
C. Thought echo.  D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	A. Persecutory delusions	
D. Illusions  E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	B. Second person auditory hallucinations	
E. Obsessional symptoms  Answer is C  338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	C. Thought echo.	
Answer is C 338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	D. Illusions	
Answer is C 338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age	E. Obsessional symptoms	
338.which of the following factors increases the risk of suicide in depression:  A. Female gender  B. Married  C. Old age		Answer is C
A. Female gender B. Married C. Old age	778 which of the following factors increases the risk of spicide in depression:	
B. Married C. Old age		
C. Old age	-	
	D. Young girls	

Answer is C

\*\*\*. Which of the following statements regarding Hypochondriasis is True?

A. Hypochondriasis is a dissociative disorder.

E. Educational grade

The patient presents with many physical symptoms related to more than one organ without a physical pathology. C. Is always a primary disorder. D.Is a preoccupation with disease that is more prevalent in middle aged people than other age groups. E. Affects females significantly more than males Very old question ,hypochondriasis means illness anxiety disorder 339. Which of the following is True of anxiety symptoms? A. Anxiety is always pathological. Low level of anxiety improves performance. C. Anxiety affects males more than females. D. Conversion disorder is associated with high anxiety level. E. Anxiety is associated with lowered autonomic arousal. Answer is B 340. Which of the following is True of phobic disorder? A. The fear is controllable. The fear is proportional to the need of the provoking stimulus. C. Response to anxiolytic medication is marked. D. Desensitization is preferred over flooding in agoraphobia E. Agoraphobia may follow a panic attack. Answer is E 341. Which of the following is True of delirium? Nocturnal improvement from delirium is characteristic. Misidentification of known people often occurs. B. Intact attention Frequently associated with visual hallucinations in clear consciousness. Patient often recalls events experienced during delirium after recovery. Answer is B 342. All the following are correct regarding acute stress disorder, EXCEPT: A. Recurrent dreams related to the event. Increased vigilance C. it affects only those who are subjected to the event. D. Avoidance of situations likely to trigger memories. E. Intrusive thoughts, feeling and perceptions related to the event. Answer is C 343.a case of paracetamol toxicity NAC

344-pt with hallucination all possible diagnosis except:

delusional disorder

345-pt with sudden cloudiness of consciousness, he was healthy before:	
Delirium	
346. chronic alcohol use man, treatment of his condition	
a-clonidine	
b-benzodiazepam	
o-oonzoudzopum	B is the answe
	b is tile aliswe
347. does not cause depression:	
A. nifedipine	
B. reserpine	Answer is A
348. Schizophrenic patient on medication, presented with weight gain, unilateral gynecomastia, cy	ytopenia. all of
the following may be the cause except:	
A. fluoxetine	
B. olanzapine C. haloperidol	
o. Amoportuot	Answer is A
349. Visual hallucination is most specific of:	
A. Schizophrenia	
B. Delirium	
	Answer is B
350. Alcoholic patient presented with diplopia, inability to stand, decrease LOC, one day after trea	ted with
(something) the most likely cause is.	
A. delirium	
B. vit B deficiency	
	Answer is B
351. Not associated with sympathetic hyperarousal:	
A. conversion disorders	
B. delirium	Answer is A
352. Patient with anorexia nervosa may have all of the following except:	Allswer is A
A. hyperkalemia B. leukopenia	
C. thrombocytosis	
	Answer is C
353.What do we use selective 5HT blockers for?	
major depression	

354. Which one is an SSR?

fluoxetine
355.Major depression patients , most common presentation is?
somatic symptoms.
356.Characteristic of schizophrenia?
auditory hallucinations
357. Paranoid delusion is a feature may be found in any of the following disorders, EXCEPT:
A. Mania
B. Psychotic depression
C. Cognitive disorder
D Schizophrenia
E. Impulse control disorder
Answer is E
358. Which of the following is true of panic disorder?
A Presenting in attacks, each lasting more than 12 hours
B. The attack is associated with fear of collapse.
c. Equally prevalent in men and women
D. Can be effectively treated by anticholinergic medication.
E. The patient usually carries high suicidal risk.
Answer is E
359. Somatization disorder may include all the following, EXCEPT:
A. Multiple physical complaints without underlying pathology
B. Dissociative fugue state
C. Symptoms related to more than one symptom.
D. Secondary depressive symptoms.
E. Pain symptoms nonresponsive to pain killers
Answer is B
360. Which of the following, on long term use, causes physical withdrawal symptoms on abrupt discontinuation?
A.Cannabis
B. Alcohol
C.Cocaine
D.Risperidone -
E.Amphetamines
Answer is I

 ${\bf 361}.$  All the following symptoms favor the diagnosis of OCD, EXCEPT:

A Hearing commands from others who are close to him. B. Excessive rumination about topics considered senseless by the patient. C. Intrusive thoughts with offensive content recognized by the patient as false. D. Recurrent doubts followed by repeated checking. E. Recurrent impulsive to act, which are recognized by the patient as unwise. Answer is A 362. Visual hallucination favors the diagnosis of one of the following conditions: A. Schizoaffective disorder B. Bipolar disorder C. Schizophrenia D. Delusional disorder E Delirium Answer is E 363. Which of the following is True of Impulse Control Disorder? A. Fugue can be a presenting feature. B. May present with trichotillomania. May present with unexplained physical symptoms. Includes malingering. Is associated with experiencing depersonalization. Answer is B 364. Which of the following is a characteristic feature of Post-Traumatic Stress Disorder? A. Involvement in conversations related to the trauma. B Recurrent distressing dreams C. Frequently visiting the place where the trauma occurred D. Lowered arousal level. E. The patient is always a survived victim of the trauma. Answer is B 365. Desensitization is effective in the treatment of: A Specific phobia B. Social phobia C. Agoraphobia D. Obsessive compulsive disorder E. Conversion disorder Answer is A 366. Which of the following medications would be the most appropriate to prescribe for a patient with Bipolar Depression? A. Haloperidol **Bromazepam** Valproic acid

D. Fluoxetine E. Risperidon Answer is C 367. Acute drug induced dystonia can be due to the use of: Procyclidine B. Alprazolam C. Propranolol D. Olanzapine E haloperidol Answer is D 368.Treated overdose of SSRI: A. OCD B. Borderline C. Panic attack Answer is A 369.In cases of recurrent manic attacks you can add what drug? carbamazepine \*\*\*\*\*DOCTOR 2018 MINI-OSCE MCQ\*\*\* MCQs: 1) Which of the following is not in the differential diagnosis of psychosis? Schizophrenia Bipolar depression Mania Unipolar depression Social Anxiety More than one of the above Answer: Social Anxiety 2) Which of the following must have delusions in order to be diagnosed? Schizophrenia Bipolar

**MDD** 

Delusional Disorder

More than one of the above

Answer: Delusional disorder

3) Which of the following is correct?

Both negative and positive symptoms must be present in active psychosis

Psychosis occurs in less than 20% of acute mania patients

In mania the mood must be elevated

None of the above

Answer: None of the above

4) Which of the following is correctly matched

Delirium -> IV Diazepam

Quetiapine-> bipolar depression

Cetalopram -> unipolar depression

More than one of the above

Answer More than one of the above (the unipolar and bipolar depression treatments are both correct)

5) Which of the following is not treated by SSRIs?

Negative symptoms of schizophrenia

Atypical depression

Melancholic features of depression

OCD

More than one of the above

Answer: Negative symptoms of schizophrenia

Q6/Case of bipolar on treatment come to the clinic with diarrhea and

tremor it's due to?

Lithium side effects

Q7/Nahilistic delusion indicate? Psychotic depression

Q8/Which of the following is true according to psychosis? Psychosis should be treated as medical health disease

Q9/ Which of the following is a disorder of perception? Illusion

- 10. Which of the following represents a disorder of thinking?
- A. Conversion
- B. Thought echo
- C. Hallucination
- D. Depersonalisation
- E. Delusions

**ANSWER IS E** 

